# SRSF3 Recombinant Rabbit Monoclonal Antibody (JE46-44)

## **Product Details**

Size	100 µL	
Species Reactivity	Human	
Host/Isotype	Rabbit / IgG	
Expression system	HEK293 cells	
Class	Recombinant Monoclonal	
Туре	Antibody	
Clone	JE46-44	
Conjugate	Unconjugated	
Immunogen	Synthetic peptide within Human SFRS3 aa 34-83	
Form	Liquid	
Concentration	1 mg/mL	
Purification	Protein A	
Storage buffer	TBS, pH 7.4, with 40% glycerol, 0.05% BSA	
Contains	0.05% sodium azide	
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles, store in dark	
RRID	AB_2848754	


Applications	Tested Dilution	Publications
Western Blot (WB)	1:500-1:2,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:100-1:500	-
Immunocytochemistry (ICC/IF)	1:50-1:200	-

# **Product Specific Information**

Positive Control: Mouse testis tissue, rat brain tissue, A431, SH-SY-5Y, SiHa, 293, Hela, HepG2, A549, MG-63.


1

# Product Images For SRSF3 Recombinant Rabbit Monoclonal Antibody (JE46-44)




### SRSF3 Antibody (MA5-34846) in WB

Western blot analysis of SFRS3 in Positive control: Lane 1: Mouse testis tissue lysate, Lane 2: Rat brain tissue lysate, Lane 3: A431 cell lysate, Lane 4: SH-SY-5Y cell lysate, Lane 5: SiHa cell lysate, Lane 6: 293 cell lysate, Lane 7: HeLa cell lysate, Lane 8: HepG2 cell lysate, Lane 9: A549 cell lysate. Samples were transferred to PVDF membrane, blocked with 5% BSA (1 hour), incubated with SFRS3 monoclonal antibody (Product # MA5-34846), at a dilution of 1:500 (5% BSA, 2 hours), followed by Goat Anti-Rabbit IgG-HRP (1 hour) with a dilution of 1: 5000.



#### SRSF3 Antibody (MA5-34846) in ICC/IF

Immunofluorescent analysis of SFRS3 in MG-63 cells (green). Samples were formalin fixed, permeabilized with 0.1% Triton X-100 in TBS (1 hour, room temperature) and blocked with 1% BSA (15 min, room temperature), incubated with SFRS3 monoclonal antibody (Product # MA5-34846) at a dilution of 1:100 (1 hour, room temperature), and followed by Alexa Fluor 488 Goat anti-Rabbit IgG and DAPI (blue) with a dilution of 1:100.



#### SRSF3 Antibody (MA5-34846) in IHC (P)

Immunohistochemistry analysis of SFRS3 in paraffin-embedded human breast cancer tissue. Samples were heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4, 20 minutes) and blocked in 5% BSA (30 min, room temperature), incubated with SFRS3 monoclonal antibody (Product # MA5-34846) at a dilution of 1:200 (30 min, room temperature), and followed by HRP conjugate, DAB and hematoxylin (mounted with DPX).

#### View more figures on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by property trained individuals. Unless otherwise stated in the Documentation, this warranty is marring to any arrow date of asimple furnished to super itemation. This warranty are not atea of aligned to asimple furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample furnished. Express OR INVERSES OR CALCEMENT OF OR NON-CONFORMING PRODUCTS DURING THE WARRANT PERIOD IS LIMITED TO REPAR, REPLACEMENT OF OR REPLAN PERIOD IS SULFERS OR CALCEMENT OF OR REPLAN PERIOD IS SULFERS OR INVERSES OR INVERE