



MAFF Polyclonal Antibody

Product Details		
Size	100 μg	
Species Reactivity	Human	
Host/Isotype	Goat / IgG	
Class	Polyclonal	
Туре	Antibody	
Conjugate	Unconjugated	
Immunogen	E. coli-derived recombinant human MafF Ser2-Ser164	
Form	Lyophilized	
Concentration	0.2 mg/mL	
Purification	Antigen affinity chromatography	
Storage buffer	PBS with 5% trehalose	
Contains	No Preservative	
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles	
RRID	AB_2576549	


Applications	Tested Dilution	Publications
Western Blot (WB)	2 μg/mL	-
Immunocytochemistry (ICC/IF)	5-15 μg/mL	-

Product Specific Information

In Western blots, less than 1% cross-reactivity with recombinant human (rh) MafG and rhMafK is observed.

Reconstitute at 0.2 mg/mL in sterile PBS.


Product Images For MAFF Polyclonal Antibody



MAFF Antibody (PA5-47552) in WB


Western blot analysis of MAFF in MDA-MB-468 human breast cancer cell line and SK-Mel-28 human malignant melanoma cell line. Samples were incubated in MAFF polyclonal antibody (Product # PA5-47552) using a dilution of 2 μ g/mL followed by a HRP-conjugated Anti-Goat IgG secondary antibody. A specific band for MafF was detected at approximately 19 kDa (as indicated). This experiment was conducted under reducing conditions.





MAFF Antibody (PA5-47552) in ICC/IF

Immunocytochemistry analysis of MAFF in immersion fixed HepG2 human hepatocellular carcinoma cell line. Samples were incubated in MAFF polyclonal antibody (Product # PA5-47552) using a dilution of 15 µg/mL for 3 hours at room temperature followed by NorthernLights[™] 557-conjugated Anti-Goat IgG Secondary Antibody (left panel, red) and counterstained with DAPI (right panel, blue). Specific staining was localized to nuclei.



MAFF Antibody (PA5-47552) in WB

Western blot analysis of MAFF in 2 ng/lane recombinant human (rh) MafF, MafG, and MafK. Samples were incubated in MAFF polyclonal antibody (Product # PA5-47552) using a dilution of 2 µg/mL followed by a HRP-conjugated Anti-Goat IgG secondary antibody. A specific band for MafF was detected at approxi-mately 19 kDa (as indicated). This experiment was conducted under reducing conditions.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, his warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample in terminished to Buyer is impressible using its impressible to the product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING IT HE WARRANTY PERSO IS LIMITED REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCTS) AT SITE LER'S SOLE OPTION. THERE IS SOLE OPTI