



Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), Super Bright™ 436, eBioscience™

| Product Details | | |
|----------------------------|-------------------------------------|--|
| Size | 100 μg | |
| Host/Isotype | Mouse / IgG1, kappa | |
| Class | Control | |
| Туре | Isotype Control | |
| Clone | P3.6.2.8.1 | |
| Conjugate | Super Bright™ 436 | |
| Excitation/Emission Max | 413/431 nm | |
| Form | Liquid | |
| Concentration | 0.2 mg/mL | |
| Purification | Affinity chromatography | |
| Storage buffer | PBS, pH 7.2, with BSA | |
| Contains | 0.09% sodium azide | |
| Storage conditions | 4° C, store in dark, DO NOT FREEZE! | |
| RRID | AB_2637446 | |

| Applications | Tested Dilution | Publications |
|-----------------------|-----------------|--------------|
| Flow Cytometry (Flow) | Assay-Dependent | - |
| Control (Ctrl) | Assay-Dependent | - |

Product Specific Information

Description: The monoclonal mouse IgG1 K immunoglobulin is useful as an isotype control.

Applications Reported: This P3.6.2.8.1 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This P3.6.2.8.1 antibody has been tested by flow cytometric analysis of normal human peripheral blood cells and mouse splenocytes. Use the isotype control at the same concentration as the experimental antibody.

Super Bright 436 can be excited with the violet laser line (405 nm) and emits at 436 nm. We recommend using a 450/50 bandpass filter, or equivalent. Please make sure that your instrument is capable of detecting this fluorochrome.

When using two or more Super Bright dye-conjugated antibodies in a staining panel, it is recommended to use Super Bright Complete Staining Buffer (Product # SB-4401) to minimize any non-specific polymer interactions. Please refer to the datasheet for Super Bright Staining Buffer for more information.

Excitation: 405 nm; Emission: 436 nm; Laser: Violet Laser

Super Bright Polymer Dyes are sold under license from Becton, Dickinson and Company.

□ 1 Reference

SHP2 inhibition enhances the anticancer effect of Osimertinib in EGFR T790M mutant lung adenocarcinoma by blocking CXCL8 loop mediated stemness Research Square (2021)

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation. Specifications and/or accompanying package inserts ("Documentation"), No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICLAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE EMBEDY FOR NON-CORPORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITACE. OR REFUND FOR THE NON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses, or any type of consumption to human or animal to the human or anima