

CD90.2 (Thy-1.2) Monoclonal Antibody (53-2.1), FITC, eBioscience™

Product Details	
Size	100 µg
Species Reactivity	Mouse
Published Species	Fish, Mouse, Human
Host/Isotype	Rat / IgG2a, kappa
Recommended Isotype Control	Rat IgG2a kappa Isotype Control (eBR2a), FITC, eBioscience™
Class	Monoclonal
Type	Antibody
Clone	53-2.1
Conjugate	FITC
Form	Liquid
Concentration	0.5 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.1% gelatin
Contains	0.09% sodium azide
Storage Conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_465154

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	0.125 µg/test	16 Publications
Immunocytochemistry (ICC)	-	1 Publication
Immunofluorescence (IF)	-	3 Publications
Immunohistochemistry (Frozen) (IHC (F))	-	1 Publication
Miscellaneous PubMed (Misc)	-	1 Publication

Product Specific Information

Description: The 53-2.1 monoclonal antibody reacts with mouse CD90.2 also known as Thy-1.2, a GPI-linked membrane molecule. CD90.2 is expressed by mouse thymocytes and mature T cells as well as neurons in CD90.2-expressing mouse strains. These strains include BALB/c, CBA, C3H, C57BL/6, C58/, SJL and others. Cells from CD90.1-expressing strains including PL and AKR do not stain with 53-2.1. CD90 is involved in regulation of adhesion and signal transduction by T cells.


Applications Reported: The 53-2.1 antibody has been reported for use in flow cytometric analysis.

Applications Tested: The 53-2.1 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.125 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Excitation: 488 nm; **Emission:** 520 nm; **Laser:** Blue Laser.

Filtration: 0.2 µm post-manufacturing filtered.

Product Images For CD90.2 (Thy-1.2) Monoclonal Antibody (53-2.1), FITC, eBioscience™



CD90.2 (Thy-1.2) Antibody (11-0902-82) in Flow

Staining of mouse splenocytes with Anti-Mouse CD90-2 (Thy-1-2) FITC (left) or PE (right). Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.

[View more figures on thermofisher.com](https://www.thermofisher.com)

Immunohistochemistry (Frozen) (1)

The American journal of pathology

Distinct developmental requirements for isolated lymphoid follicle formation in the small and large intestine: RANKL is essential only in the small intestine.

"11-0902 was used in Immunofluorescence to study the role of RANKL in the development of cryptopatches and isolated lymphoid follicles within intestinal lymphoid tissues."

Authors: Knoop KA,Butler BR,Kumar N,Newberry RD,Williams IR

Species
Mouse

Dilution
Not Cited

Year
2011

Immunocytochemistry (1)

Proceedings of the National Academy of Sciences of the United States of America

Dysregulated YAP1/TAZ and TGF- signaling mediate hepatocarcinogenesis in Mob1a/1b-deficient mice.

"11-0902 was used in Immunofluorescence to study the molecular mechanisms underpinning the pathology of Mob1a/1b-deficient mice."

Authors: Nishio M,Sugimachi K,Goto H,Wang J,Morikawa T,Miyachi Y,Takano Y,Hikasa H,Itoh T,Suzuki SO,Kurihara H,Aishima S,Leask A,Sasaki T,Nakano T,Nishina H,Nishikawa Y,Sekido Y,Nakao K,Shin-Ya K,Mimori K,Suzuki A

Species
Mouse

Dilution
Not Cited

Year
2016

Immunofluorescence (3)

Nature

Single-cell transcriptomics reconstructs fate conversion from fibroblast to cardiomyocyte.

"Published figure using CD90.2 (Thy-1.2) monoclonal antibody (Product # 11-0902-82) in Immunofluorescence"

Authors: Liu Z,Wang L,Welch JD,Ma H,Zhou Y,Vaseghi HR,Yu S,Wall JB,Alimohamadi S,Zheng M,Yin C,Shen W,Prins JF,Liu J,Qian L

Species
Not Applicable

Dilution
Not Cited

Year
2017

The American journal of pathology

Distinct developmental requirements for isolated lymphoid follicle formation in the small and large intestine: RANKL is essential only in the small intestine.

"11-0902 was used in Immunofluorescence to study the role of RANKL in the development of cryptopatches and isolated lymphoid follicles within intestinal lymphoid tissues."

Authors: Knoop KA,Butler BR,Kumar N,Newberry RD,Williams IR

Species
Mouse

Dilution
Not Cited

Year
2011

[View more IF references on thermofisher.com](#)

More applications with references on thermofisher.com

Flow (16)

Misc (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.