



IL-17F Monoclonal Antibody (SHLR17), PE, eBioscience™

Product Details	
Size	100 Tests
Species Reactivity	Human
Published Species	Human
Host/Isotype	Rat / IgG1, kappa
Recommended Isotype Control	Rat IgG1 kappa Isotype Control (eBRG1), PE, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	SHLR17
Conjugate	PE
Excitation/Emission Max	565/576 nm
Form	Liquid
Concentration	5 μL/Test
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.2% BSA
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2572654

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	5 μL (0.5 μg)/test	1 Publication

Product Specific Information

Description: The SHLR17 monoclonal antibody reacts with human interleukin (IL)-17F. IL-17F is a homodimeric, 34 kDa member of the IL-17 family, which includes IL-17(A), B, C, D, E (IL-25). Within this family, IL-17A and IL-17F are produced preferentially by T helper 17 (Th17) cells, with other members produced more widely. IL-17F and IL-17A have many overlapping functions, but knockout studies have shown each to have independent functions as well. In vitro cultured Th17 cells have been shown to express IL-17F and IL-17A homodimers and IL-17A/F heterodimers depending on the culture conditions and their differentiation state. A heterodimer of IL-17RA and IL-17RC has been shown to bind IL-17F, resulting in target cell-secretion of pro-inflammatory cytokines and chemokines, as well as neutrophil recruitment.


Applications Reported: This SHLR17 antibody has been reported for use in intracellular staining followed by flow cytometric analysis.


Applications Tested: This SHLR17 antibody has been pre-titrated and tested by intracellular staining followed by flow cytometric analysis of TH17-polarized normal human peripheral blood cells. This can be used at 5 µL (0.5 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

Excitation: 488-561 nm; Emission: 578 nm; Laser: Blue Laser, Green Laser, Yellow-Green Laser.

Filtration: 0.2 µm post-manufacturing filtered.

Product Images For IL-17F Monoclonal Antibody (SHLR17), PE, eBioscience™





IL-17F Antibody (12-7169-42) in Flow

Intracellular staining of CD4-enriched Th17-polarized, (using Human IL-23 Recombinant Protein (Product # 14-8239-63) normal human peripheral blood cells after 5-hour stimulation with Cell Stimulation Cocktail (plus protein transport inhibitors) (Product # 00-4975-03) with Anti-Human IL-17A Alexa Fluor® 647 and Mouse IgG1 K Isotype Control PE (Product # 12-4714-81) (left) or Anti-Human IL-17F PE (right). Cells were fixed and premeabilized using Intracellular Fixation & Permeabilization Buffer Set (Product # 88-8824-00). Total viable cells were used for analysis.

□ 1 Reference

Flow Cytometry (1)

Journal of immunology (Baltimore, Md.: 1950)

Dimethyl Fumarate Selectively Reduces Memory T Cells and Shifts the Balance between Th1/Th17 and Th2 in Multiple Sclerosis Patients.

"12-7169 was used in Flow cytometry/Cell sorting to study the therapeutic effects of dimethyl fumarate and its side effect of progressive multifocal leukoencephalopathy."

Authors: Wu Q, Wang Q, Mao G, Dowling CA, Lundy SK, Mao-Draayer Y

Year 2017

Species Human

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production obcumentation, specifications and/or accompanying package insents ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is imited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anynous other than the Buyer. Any mode of sample in the Product is will conform to such model or sample incomplete the property of the property of