

# TRA-1-60 (Podocalyxin) Monoclonal Antibody (TRA-1-60), PE, eBioscience™

## Product Details

|                         |                                     |
|-------------------------|-------------------------------------|
| Size                    | 100 µg                              |
| Species Reactivity      | Human                               |
| Published Species       | Human, Mouse                        |
| Host/Isotype            | Mouse / IgM                         |
| Class                   | Monoclonal                          |
| Type                    | Antibody                            |
| Clone                   | TRA-1-60                            |
| Conjugate               | PE                                  |
| Excitation/Emission Max | 565/576 nm                          |
| Form                    | Liquid                              |
| Concentration           | 0.2 mg/mL                           |
| Purification            | Affinity chromatography             |
| Storage buffer          | PBS, pH 7.2                         |
| Contains                | 0.09% sodium azide                  |
| Storage conditions      | 4° C, store in dark, DO NOT FREEZE! |
| RRID                    | AB_891602                           |

| Applications                 | Tested Dilution | Publications    |
|------------------------------|-----------------|-----------------|
| Immunocytochemistry (ICC/IF) | -               | 1 Publication   |
| Flow Cytometry (Flow)        | 1 µg/test       | 10 Publications |

## Product Specific Information

**Description:** The TRA-1-60 antibody recognizes a protein expressed on undifferentiated human embryonic stem cells (ES), embryonal carcinoma cells (EC), and embryonic germ cells (EG). Like other stem cell specific markers, the epitope recognized by the TRA-1-60 antibody is lost upon cell differentiation. Contrary to early reports that the TRA-1-60 epitope can be destroyed by neuraminidase digestion, new studies have shown that TRA-1-60 recognizes a neurominidase resistant antigen. The TRA-1-60 antibody is known to specifically recognize a carbohydrate epitope on a keratan sulfated glycoprotein recently identified as podocalyxin, a member of the CD34-related family of sialomucins. Podocalyxin is a transmembrane glycoprotein originally identified on epithelial glomerular cells known as podocytes, and the protein has also been implicated in the development of aggressiveness in a variety of cancers, including breast and prostate cancer.


**Applications Reported:** This TRA-1-60 antibody has been reported for use in flow cytometric analysis.

**Applications Tested:** This TRA-1-60 antibody has been tested by flow cytometric analysis of the human embryonal carcinoma (EC) line 2102Ep. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

**Excitation:** 488-561 nm; **Emission:** 578 nm; **Laser:** Blue Laser, Green Laser, Yellow-Green Laser.

**Filtration:** 0.2 µm post-manufacturing filtered.

Product Images For TRA-1-60 (Podocalyxin) Monoclonal Antibody (TRA-1-60), PE, eBioscience™



**TRA-1-60 (Podocalyxin) Antibody (12-8863-82) in Flow**  
Staining of 2102Ep human embryonal carcinoma cell line with 0.5 µg of Mouse IgM Isotype Control PE (Product # 12-4752) (open histogram) or 0.5 µg of Anti-Human TRA-1-60 PE (filled histogram). Total viable cells were used for analysis.

View more figures on [thermofisher.com](https://thermofisher.com)

11 References

Immunocytochemistry (1)

|   |                  |
|---|------------------|
| Cell stem cell  | Year<br>2017     |
| <b>Analysis of Transcriptional Variability in a Large Human iPSC Library Reveals Genetic and Non-genetic Determinants of Heterogeneity.</b>   | Species<br>Human |
| "12-8863 was used in Immunocytochemistry to examine the sources of gene expression variability in 317 human induced pluripotent stem cell lines from 101 individuals."  |                  |
| Authors: Carcamo-Orive I,Hoffman GE,Cundiff P,Beckmann ND,D'Souza SL,Knowles JW,Patel A,Papatsenko D,Abbasi F,Reaven GM,Whalen S,Lee P,Shahbazi M,Henrion MYR,Zhu K,Wang S,Roussos P,Schadt EE,Pandey G,Chang R,Quertermous T,Lemischka I |                  |

Flow Cytometry (10)

|   |                   |
|---|-------------------|
| Journal of cellular physiology  | Year<br>2021      |
| <b>Characterization of intracellular buffering power in human induced pluripotent stem cells and the loss of pluripotency is delayed by acidic stimulation and increase of NHE1 activity.</b>   | Species<br>Human  |
| "12-8863-82 was used in Flow cytometry/Cell sorting to establish a quantitative model of passive during differentiation and demonstrate that maintenance of NHE1 at a higher level is of critical importance for pluripotency retention in hiPSCs." |                   |
| Authors: Lee SP,Chao SC,Chou MF,Huang SF,Dai NT,Wu GJ,Tsai CS,Loh SH,Tsai YT  |                   |
|   | Dilution<br>1:100 |

|   |                  |
|---|------------------|
| Cell reports  | Year<br>2019     |
| <b>Ultra-High-Frequency Reprogramming of Individual Long-Term Hematopoietic Stem Cells Yields Low Somatic Variant Induced Pluripotent Stem Cells.</b> | Species<br>Human |
| "12-8863-82 was used in Flow Cytometry to suggest LT-HSCs may become the preferred cell source for the production of clinical-grade iPSCs."           |                  |
| Authors: Wang K,Guzman AK,Yan Z,Zhang S,Hu MY,Hamaneh MB,Yu YK,Tolu S,Zhang J,Kanavy HE,Ye K,Bartholdy B,Bouhassira EE                                |                  |

View more Flow references on [thermofisher.com](https://thermofisher.com)

More applications with references on [thermofisher.com](https://thermofisher.com)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.