




PCNA Monoclonal Antibody (PC10), Biotin

Product Details	
Size	100 μg
Species Reactivity	Human, Insect, Mouse, Non-human primate, Rat, Yeast
Published Species	Sheep, Human, Mouse
Host/Isotype	Mouse / IgG2a
Class	Monoclonal
Туре	Antibody
Clone	PC10
Conjugate	Biotin
Immunogen	Rat recombinant PCNA polypeptides.
Form	Liquid
Concentration	0.2 mg/mL
Purification	purified
Storage buffer	PBS, pH 7.4, with 50% glycerol, 1% BSA
Contains	0.1% sodium azide
Storage conditions	4° C
RRID	AB_2533017


Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000	2 Publications
Immunohistochemistry (IHC)	-	5 Publications
Immunohistochemistry (Paraffin) (IHC (P))	1:10-1:100	4 Publications
Immunocytochemistry (ICC/IF)	-	2 Publications
Miscellaneous PubMed (Misc)	-	1 Publication

Product Images For PCNA Monoclonal Antibody (PC10), Biotin




PCNA Antibody (13-3940)

Antibody specificity was demonstrated by siRNA mediated knockdown of target protein. HeLa cells were transfected with PCNA siRNA and loss of signal was observed in Western Blot using Anti-PCNA monoclonal Antibody (Product # 13-3940). {KD}



PCNA Antibody (13-3940) in WB

Western blot analysis of PCNA was performed by loading 20 µg of HeLa (lane1), Raji (lane2), MCF-7 (lane3), Jurkat (lane4), HEK-293 (lane5), NIH/3T3 (lane6), MDA-MB-231 (lane7) cell lysate using Novex® NuPAGE® 4-12 % Bis-Tris gel (Product # NP0322BOX), XCell SureLock™ Electrophoresis System (Product # El0002), Novex® Sharp Pre-Stained Protein Standard (LC5800), and iBlot® Dry Blotting System (IB21001). Proteins were transferred to a nitrocellulose membrane and blocked with 5 % skim milk for 1 hour at room temperature. PCNA was detected at ~ 34 kDa using PCNA, Biotin conjugate Mouse Monoclonal Antibody (Product # 13-3940) at 1:1000 dilution in 5 % skim milk at 4° C overnight on a rocking platform. Streptavidin - HRP (Product # SA1007) at 1: 2000 dilution was used and chemiluminescent detection was performed using Pierce™ ECL Western Blotting Substrate (Product # 32106).



PCNA Antibody (13-3940) in WB

Knockdown of PCNA was achieved by transfecting HeLa cells with PCNA specific validated siRNAs (Silencer® select Product # S10135, S10133). Western blot analysis (Fig. a) was performed using membrane enriched extracts from the PCNA knockdown cells (lane 3), non-specific scrambled siRNA transfected cells (lane 2) and untransfected cells (lane 1). The blots were probed with PCNA Monoclonal Antibody, Biotin (Product # 13-3940, 1:1000 dilution) and Streptavidin-HRP (Product # N200, 1:10,000 dilution). Densitometric analysis of this western blot is shown in histogram (Fig. b). Decrease in signal upon siRNA mediated knock down confirms that antibody is specific to PCNA .

View more figures on thermofisher.com

□ 14 References

Western Blot (2)

Journal of orthopaedic surgery and research

Expression of STOML2 promotes proliferation and glycolysis of multiple myeloma cells via upregulating PAI-1.

"13-3940 was used in Western Blotting to demonstrate that STOML2 silencing inhibits cell proliferation and glycolysis through downregulation of PAI-1 expression, suggesting a new therapeutic target for MM."

Authors: Zhang H, Wu G, Feng J, Lu X, Liu P

Year 2021

Species Human

Dilution 1:2000

Cancer research

The 104-123 amino acid sequence of the beta-domain of von Hippel-Lindau gene product is sufficient to inhibit renal tumor growth and invasion.

Authors: Datta K,Sundberg C,Karumanchi SA,Mukhopadhyay D

Year 2001

Species Human

Immunohistochemistry (5)

PLoS pathogens

HMGB1 amplifies ILC2-induced type-2 inflammation and airway smooth muscle remodelling.

"13-3940 was used in Immunohistochemistry to investigate the processes that mediate tissue damage-induced type-2 inflammation and the origins of airway remodelling, using a preclinical mouse model of viral bronchiolitis."

Authors: Loh Z,Simpson J,Ullah A,Zhang V,Gan WJ,Lynch JP,Werder RB,Sikder AA,Lane K,Sim CB,Porrello E, Mazzone SB,Sly PD,Steptoe RJ,Spann KM,Sukkar MB,Upham JW,Phipps S

Year 2020

Species Mouse

Cell reports

JNK Promotes Epithelial Cell Anoikis by Transcriptional and Posttranslational Regulation of BH3-Only Proteins.

"13-3940 was used in Immunohistochemistry-immunofluorescence to study the role of the cJUN NH2-terminal kinase signalling pathway in anoikis."

Authors: Girnius N, Davis RJ

Year 2017

Species Mouse

Dilution 1:50

View more IHC references on thermofisher.com

More applications with references on thermofisher.com

IHC (P) (4) ICC/IF (2)

Misc (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.