



GFAP Monoclonal Antibody (GA5), eBioscience™

Product Details		
Size	100 μg	
Species Reactivity	Chicken, Human, Mouse, Pig, Rabbit, Rat	
Published Species	Rat, Human, Mouse	
Host/Isotype	Mouse / IgG1	
Class	Monoclonal	
Туре	Antibody	
Clone	GA5	
Conjugate	Unconjugated	
Form	Liquid	
Concentration	0.5 mg/mL	
Purification	Affinity chromatography	
Storage buffer	PBS, pH 7.2	
Contains	0.09% sodium azide	
Storage conditions	4° C	
RRID	AB_10598206	

Applications	Tested Dilution	Publications
Western Blot (WB)	0.5-5 μg/mL	3 Publications
Immunohistochemistry (IHC)	-	19 Publications
Immunohistochemistry (Paraffin) (IHC (P))	5 μg/mL	1 Publication
Immunohistochemistry (Frozen) (IHC (F))	-	4 Publications
Immunocytochemistry (ICC/IF)	5 μg/mL	8 Publications
Flow Cytometry (Flow)	Assay-Dependent	4 Publications

Product Specific Information

Description: This GA5 monoclonal antibody reacts with human, mouse, rat, chicken, rabbit, and pig glial fibrillary acidic protein (GFAP). This 49-kDa type III intermediate filament protein is expressed in neural tissues and distinguishes astrocytes from other glial cells during central nervous system development. Three alternative splice variants of GFAP exist; however, alpha-GFAP is the predominant form expressed in astrocytes. GFAP can co-assemble with vimentin and nestin in astrocytes, but such associations are not required for assembly. Like other intermediate filaments, GFAP assembly is dependent on phosphorylation and dephosphorylation of the N-terminal domain. Studies have demonstrated that mutations in the GFAP gene lead to Alexander disease. Moreover, GFAP has also been shown to be overexpressed in certain glial-derived tumors.

Applications Reported: This GA5 antibody has been reported for use in immunocytochemistry, western blotting, immunohistochemical staining of formalin-fixed paraffin embedded tissue sections, and flow cytometric analysis.


Applications Tested: This GA5 antibody has been tested by immunoblotting of lysate prepared from mouse brain, immunocytochemistry of fixed and permeabilized C6 cells, and immunohistochemistry of FFPE human tissue using low pH antigen retrieval. This can be used at less than or equal to 5 µg/mL.

Purity: Greater than 90%, as determined by SDS-PAGE.

Aggregation: Less than 10%, as determined by HPLC.


Filtration: 0.2 µm post-manufacturing filtered.

Product Images For GFAP Monoclonal Antibody (GA5), eBioscience™




GFAP Antibody (14-9892-82)

Antibody specificity was demonstrated by detection of differential basal expression of the target across tissue tested owing to their inherent genetic constitution. Relative expression of GFAP was observed in Mouse brain, Rat brain in comparison to Mouse liver and Rat liver using Anti-GFAP Monoclonal Antibody (GA5), eBioscienceTM(Product # 14-9892-82) in Western Blot. {RE}



GFAP Antibody (14-9892-82) in WB

Western blot was performed using Anti-GFAP Monoclonal Antibody (GA5), eBioscienceTM(Product # 14-9892-82) and a 50 kDa band corresponding to GFAP was observed across tissues tested except Mouse and Rat liver. Whole cell extracts (30 μg lysate) of Mouse brain (Lane 1), Rat brain (Lane 2), Mouse liver (Lane 3) and Rat liver (Lane 4) were electrophoresed using Novex® NuPAGE® 4-12 % Bis-Tris gel (Product # NP0321BOX). Resolved proteins were then transferred onto a nitrocellulose membrane (Product # IB23001) by iBlot® 2 Dry Blotting System (Product # IB21001). The blot was probed with the primary antibody (0.5 μg/ml) and detected by chemiluminescence Goat anti-Mouse IgG (H+L), SuperclonalTM Recombinant Secondary Antibody, HRP (Product # A28177, 1:4000 dilution) using the iBright FL 1000 (Product # A32752). Chemiluminescent detection was performed using Novex® ECL Chemiluminescent Substrate Reagent Kit (Product # WP20005).



GFAP Antibody (14-9892-82) in WB

Western blot was performed using Anti-GFAP Monoclonal Antibody (GA5), eBioscience™(Product # 14-9892-80) and a 50 kDa band corresponding to GFAP was observed across tissues tested except Mouse and Rat liver. Whole cell extracts (30 µg lysate) of Mouse brain (Lane 1), Rat brain (Lane 2), Mouse liver (Lane 3) and Rat liver (Lane 4) were electrophoresed using Novex® NuPAGE® 4-12 % Bis-Tris gel (Product # NP0321BOX). Resolved proteins were then transferred onto a nitrocellulose membrane (Product # IB23001) by iBlot® 2 Dry Blotting System (Product # IB21001). The blot was probed with the primary antibody (0.5 ug/ml) and detected by chemiluminescence Goat anti-Mouse IgG (H+L), Superclonal™ Recombinant Secondary Antibody, HRP (Product # A28177, 1:4000 dilution) using the iBright FL 1000 (Product # A32752). Chemiluminescent detection was performed using Novex® ECL Chemiluminescent Substrate Reagent Kit (Product # WP20005).

View more figures on thermofisher.com

□ 39 References

Western Blot (3)

Frontiers in psychiatry

Plasma Levels of Neuron- and Astrocyte-Derived Exosomal Amyloid Beta1-42, Amyloid Beta1-40, and Phosphorylated Tau Levels in Schizophrenia Patients and Non-psychiatric Comparison Subjects: Relationships With Cognitive Functioning and Psychopathology.

"Published figure using GFAP monoclonal antibody (Product # 14-9892-82) in Western Blot"

Authors: Lee EE, Winston-Gray C, Barlow JW, Rissman RA, Jeste DV

Year 2021

Species Human

Dilution 1:1000

Neural regeneration research

Muscovite nanoparticles mitigate neuropathic pain by modulating the inflammatory response and neuroglial activation in the spinal cord.

"Published figure using GFAP monoclonal antibody (Product # 14-9892-82) in Immunohistochemistry (Frozen)" Authors: Oh JY,Hwang TY,Jang JH,Park JY,Ryu Y,Lee H,Park HJ

Year 2020

Species Mouse

Dilution 1:500

View more WB references on thermofisher.com

Immunohistochemistry (19)

International journal of molecular sciences

Inhibition of Calpain Attenuates Degeneration of Substantia Nigra Neurons in the Rotenone Rat Model of Parkinson's Disease.

"14-9892-82 was used in Immunohistochemistry to find that both isoforms are activated in a nigrostriatal pathway with increased phosphorylated synuclein following the administration of rotenone in Lewis rats, but calpain isoforms played different roles in neuronal survival."

Authors: Zaman V, Drasites KP, Myatich A, Shams R, Shields DC, Matzelle D, Haque A, Banik NL

Year 2022

Species Rat

Frontiers in immunology

Glial receptor PLXNB2 regulates schizophrenia-related stress perception *via* the amygdala.

"Published figure using GFAP monoclonal antibody (Product # 14-9892-82) in Immunohistochemistry"

Authors: Xuan FL,Yan L,Li Y,Fan F,Deng H,Gou M,Chithanathan K,Heinla I,Yuan L,Seppa K,Zharkovsky A,Kalda A, Hong LE,Hu GF,Tan Y,Tian L

Year 2022

View more IHC references on thermofisher.com

More applications with references on thermofisher.com

IHC (P) (1) IHC (F) (4) ICC/IF (8) Flow (4)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"), No claim of suitability for use in applications regulated by EDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is immitted to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample increase in merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPERSS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY END AND PRODUCTS OF A STATE OF A STA