Performance guarenteed

CD44 Monoclonal Antibody (IM7), APC, eBioscience™

Product Details

FIGUUCI Details	
Size	100 µg
Species Reactivity	Human, Mouse
Published Species	Dog, Fruit fly, Non-human primate, Bacteria, Mouse, Human
Host/Isotype	Rat / IgG2b, kappa
Recommended Isotype Control	Rat IgG2b kappa Isotype Control (eB149/10H5), APC, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	IM7
Conjugate	APC
Excitation/Emission Max	651/660 nm
Form	Liquid
Concentration	0.2 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_469390

Applications	Tested Dilution	Publications
Immunocytochemistry (ICC/IF)	-	3 Publications
Flow Cytometry (Flow)	0.06 μg/test	282 Publications
Miscellaneous PubMed (Misc)	-	1 Publication

Product Specific Information

Description: The IM7 monoclonal antibody reacts with all isoforms of mouse CD44 (Pgp-1). CD44 is expressed by hematopoietic and non-hematopoietic cells. Bone marrow myeloid cells and memory T cells highly express this antigen and peripheral B and T cells can upregulate the expression of CD44. CD44 functions as an adhesion molecule through its binding to hyaluronate, an extracellular matrix component.

Applications Reported: The IM7 antibody has been reported for use in flow cytometric analysis.


Applications Tested: The IM7 antibody has been tested by flow cytometric analysis of mouse bone marrow cells and splenocytes. This can be used at less than or equal to 0.06 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Excitation: 633-647 nm; Emission: 660 nm; Laser: Red Laser.

Filtration: 0.2 µm post-manufacturing filtered.

1

Product Images For CD44 Monoclonal Antibody (IM7), APC, eBioscience™



CD44 Antibody (17-0441-82) in Flow

Staining of C57BL/6 splenocytes with staining buffer (autofluorescence) (open histogram) or 0.03 µg of Anti-Human/Mouse CD44 APC (filled histogram). Total viable cells were used for analysis.

View more figures on thermofisher.com

286 References

Immunocytochemistry (3)

JBMR plus Year 2022 Parathyroid Hormone-Related Protein Inhibition Blocks Triple-Negative Breast Cancer Expansion in Bone Through Epithelial to Mesenchymal Species Human **Transition Reversal.** "17-0441-82 was used in Immunocytochemistry-immunoflourescence to conclude that PTHrP is a targetable EMT molecular driver and suggest that its pharmacological blockade can provide a potential therapeutic approach against established TNBC-derived skeletal lesions. Authors: Li J,Camirand A,Zakikhani M,Sellin K,Guo Y,Luan X,Mihalcioiu C,Kremer R **PloS one** Year 2020 Graphene-based 2D constructs for enhanced fibroblast support. Species "17-0441 was used in Immunocytochemistry-immunoflourescence to study pristine graphene and two of its oxygenfunctionalised derivatives-high and low-oxygen graphene films-as potential substrates for skin cell proliferation and Human differentiation. Authors: Safina I, Bourdo SE, Algazali KM, Kannarpady G, Watanabe F, Vang KB, Biris AS

View more ICC/IF references on thermofisher.com

Flow Cytometry (282)

Bioactive materials A platinum@polymer-catechol nanobraker enables radio- immunotherapy for crippling melanoma tumorigenesis, angiogenesis,	Year 2023
and radioresistance.	
"Published figure using CD44 monoclonal antibody (Product # 17-0441-82) in Flow Cytometry"	
Authors: Li W,Yan J,Tian H,Li B,Wang G,Sang W,Zhang Z,Zhang X,Dai Y	
Nature immunology	Year
IFN binding to extracellular matrix prevents fatal systemic toxicity.	2023
"17-0441-82 was used in Flow cytometry/Cell sorting to conclude that local retention of IFN is a pivotal mechanism to protect the organism from systemic toxicity during prolonged immune stimulation."	Species Mouse

protect the organism from systemic toxicity during prolonged immune stimulation." Authors: Kemna J,Gout E,Daniau L,Lao J,Weißert K,Ammann S,Kühn R,Richter M,Molenda C,Sporbert A,Zocholl D, Klopfleisch R,Lortat-Jacob H,Aichele P,Kammertoens T,Blankenstein T

View more Flow references on thermofisher.com

More applications with references on thermofisher.com

Misc (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No OTHER WARRANTIES, EXERSIS OR IMPLED, ARERSIS OR INPERION ECONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR REPUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR REPUND FOR THE RODUCTS AS THE RESULT OF () ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, () MINSUE, FAULT OR NEGLIGENCE OF OR BY BUYER, () USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR ((V) IMPROPER STORAGE AND HANDLING OFT THE PRODUCTS. Not in the det on the roduct or in the documentation accompanying the Product, is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic u