



# CD4 Monoclonal Antibody (RPA-T4), PE-Cyanine7, eBioscience™

| Product Details                |  |
|--------------------------------|--|
| Size                           | 100 Tests  |
| Species Reactivity             | Human  |
| Published Species              | Human, Mouse   |
| Host/Isotype                   | Mouse / IgG1, kappa  |
| Recommended Isotype<br>Control | Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), PE-Cyanine7, eBioscience™ |
| Class                          | Monoclonal   |
| Туре                           | Antibody   |
| Clone                          | RPA-T4   |
| Conjugate                      | PE-Cyanine7  |
| Excitation/Emission<br>Max     | 569/780 nm   |
| Form                           | Liquid   |
| Concentration                  | 5 μL/Test  |
| Purification                   | Affinity chromatography  |
| Storage buffer                 | PBS, pH 7.2, with 0.2% BSA   |
| Contains                       | 0.09% sodium azide   |
| Storage conditions             | 4° C, store in dark, DO NOT FREEZE!                                      |
| RRID                           | AB_1659695   |
|                                |  |

| Applications          | Tested Dilution    | Publications    |
|-----------------------|--------------------|-----------------|
| Flow Cytometry (Flow) | 5 μL (0.5 μg)/test | 60 Publications |

#### **Product Specific Information**

Description: The RPA-T4 monoclonal antibody reacts with human CD4, a 59 kDa cell surface receptor expressed by a majority of thymocytes, subpopulation of mature T cells (T-helper cells) and in low levels on monocytes. CD4 is a receptor for the human immunodeficiency virus (HIV). RPA-T4 blocks HIV binding and mixed lymphocyte reaction. The RPA-T4 antibody recognizes a different epitope than the OKT4 monoclonal antibody, and these antibodies do not cross-block binding to each other's respective epitopes.

Applications Reported: The RPA-T4 antibody has been reported for use in flow cytometric analysis.


Applications Tested: This RPA-T4 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5  $\mu$ L (0.5  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells /test.

Light sensitivity: This tandem dye is sensitive photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100  $\mu$ L cell sample + 100  $\mu$ L IC Fixation Buffer) or 1-step Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency /compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

Excitation: 488-561 nm; Emission: 775 nm; Laser: Blue Laser, Green Laser, Yellow-Green Laser.

### Product Images For CD4 Monoclonal Antibody (RPA-T4), PE-Cyanine7, eBioscience™



### CD4 Antibody (25-0049-42) in Flow

Staining of normal human peripheral blood cells with staining buffer (autofluorescence) (open histogram) or Anti-Human CD4 PE-Cyanine7 (filled histogram). Cells in the lymphocyte gate were used for analysis.

# View more figures on thermofisher.com

#### **□ 60 References**

# Flow Cytometry (60)

JCI insight

# Rapamycin improves Graves' orbitopathy by suppressing CD4+cytotoxic T lymphocytes.

"Published figure using CD4 monoclonal antibody (Product # 25-0049-42) in Flow Cytometry"

 $\label{thm:model} Authors: Zhang M, Chong KK, Chen ZY, Guo H, Liu YF, Kang YY, Li YJ, Shi TT, Lai KK, He MQ, Ye K, Kahaly GJ, Shi BY, Wang Y$ 

**Year** 2023

#### Frontiers in immunology

# Single-cell RNA sequencing reveals the molecular features of peripheral blood immune cells in children, adults and centenarians.

"Published figure using CD4 monoclonal antibody (Product # 25-0049-42) in Flow Cytometry"

Authors: Zhong J,Ding R,Jiang H,Li L,Wan J,Feng X,Chen M,Peng L,Li X,Lin J,Yang H,Wang M,Li Q,Chen Q

**Year** 2023

View more Flow references on thermofisher.com

#### More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation,"), No claim of subticibility for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is imited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general pand quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCTS; AT SELLER'S SOLE OPTION. THE SELLER'S SOLE OPTION. THE PROPUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, are vivo or in vivo therapeutic uses, and the vivo therapeutic uses, are vivo or in vivo therapeutic use