# CD44 Monoclonal Antibody (IM7), PE-Cyanine7, eBioscience™

## Product Details

FIGUUCI Details	
Size	100 µg
Species Reactivity	Human, Mouse
Published Species	Fruit fly, Non-human primate, Mouse, Human
Host/Isotype	Rat / IgG2b, kappa
Recommended Isotype Control	Rat IgG2b kappa Isotype Control (eB149/10H5), PE-Cyanine7, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	IM7
Conjugate	PE-Cyanine7
Excitation/Emission Max	569/780 nm
Form	Liquid
Concentration	0.2 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_469623

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	0.125 µg/test	219 Publications

## **Product Specific Information**

Description: The IM7 monoclonal antibody reacts with all isoforms of mouse CD44 (Pgp-1). CD44 is expressed by hematopoietic and non-hematopoietic cells. Bone marrow myeloid cells and memory T cells highly express this antigen and peripheral B and T cells can upregulate the expression of CD44. CD44 functions as an adhesion molecule through its binding to hyaluronate, an extracellular matrix component.

Applications Reported: This IM7 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This IM7 antibody has been tested by flow cytometric analysis of mouse bone marrow cells and splenocytes. This can be used at less than or equal to 0.125  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Light sensitivity: This tandem dye is sensitive photo-induced oxidation. Please protect this vial and stained samples from light.


Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100 µL cell sample + 100 µL IC Fixation Buffer) or 1-step Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency /compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

1

Excitation: 488-561 nm; Emission: 775 nm; Laser: Blue Laser, Green Laser, Yellow-Green Laser.

Filtration: 0.2 µm post-manufacturing filtered.

# Product Images For CD44 Monoclonal Antibody (IM7), PE-Cyanine7, eBioscience™



#### CD44 Antibody (25-0441-82) in Flow

Staining of C57BL/6 splenocytes with 0.06 µg of Rat IgG2b K Isotype Control PE-Cyanine7 (Product # 25-4031-82) (open histogram) or 0.06 µg of Anti-Human /Mouse CD44 PE-Cyanine7 (filled histogram). Cells in the lymphocyte gate were used for analysis.

## View more figures on thermofisher.com

## 219 References

#### Flow Cytometry (219)

Bioactive materials A platinum@polymer-catechol nanobraker enables radio- immunotherapy for crippling melanoma tumorigenesis, angiogenesis, and radioresistance. "Published figure using CD44 monoclonal antibody (Product # 25-0441-82) in Flow Cytometry" Authors: Li W,Yan J,Tian H,Li B,Wang G,Sang W,Zhang Z,Zhang X,Dai Y	<b>Year</b> 2023
Frontiers in immunology TRAF6 controls T cell homeostasis by maintaining the equilibrium of MALT1 scaffolding and protease functions. "Published figure using CD44 monoclonal antibody (Product # 25-0441-82) in Flow Cytometry" Authors: O'Neill TJ,Gewies A,Seeholzer T,Krappmann D	<b>Year</b> 2023

View more Flow references on thermofisher.com

# More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and general to perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, this warranty is specifications and for accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is vailed only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is imited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTES, EXPRESS OR INNELD, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTES OF MERCHANTABILITY, ITNESS FOR ANY PARTICLAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMENT FOR NON-CONFORMING PRODUCTS DURING THE WARRANTE PREVIDO IS LIMITED TO REPAR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS SOLE OPTION. THERE IS NO OBLICATION TO REPAR, REPLACE OR REFUND FOR THE RON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLICENCE O OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic

2