

CD3 Monoclonal Antibody (SK7), PE-Cyanine5.5, eBioscience™

Product Details	
Size	100 Tests
Species Reactivity	Chimpanzee, Human
Published Species	Human
Host/Isotype	Mouse / IgG1, kappa
Recommended Isotype Control	Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), PE-Cyanine5.5, eBioscience™
Class	Monoclonal
Type	Antibody
Clone	SK7
Conjugate	PE-Cyanine5.5
Excitation/Emission Max	564/692 nm
Form	Liquid
Concentration	5 µL/Test
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.2% BSA
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_11220085

Applications	Tested Dilution	Publications
Immunohistochemistry (Frozen) (IHC (F))	-	1 Publication
Immunocytochemistry (ICC/IF)	-	1 Publication
Flow Cytometry (Flow)	5 µL (0.125 µg)/test	15 Publications

Product Specific Information

Description: The SK7 monoclonal antibody reacts with human and chimpanzee CD3e, a 20 kDa subunit of the TCR complex. Along with the other CD3 subunits gamma and delta, the epsilon chain is required for proper assembly, trafficking and surface expression of the TCR complex. CD3 is expressed by thymocytes in a developmentally regulated manner and by all mature T cells. The SK7 and UCHT1 monoclonal antibodies cross-block binding, suggesting recognition of overlapping epitope. In contrast, clones OKT3 and SK7 see different epitopes.

The antibody SK7 recognizes chimpanzee CD3.

Applications Reported: This SK7 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This SK7 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 µL (0.125 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

Light sensitivity: This tandem dye is sensitive photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100 µL cell sample + 100 µL IC Fixation Buffer) or 1-step


Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency /compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

Excitation: 488-561 nm; Emission: 695 nm; Laser: Blue Laser, Green Laser, Yellow-Green Laser

Product Images For CD3 Monoclonal Antibody (SK7), PE-Cyanine5.5, eBioscience™


CD3 Antibody (35-0036-42)

Staining of human peripheral blood cells. As expected based on known relative expression patterns, CD3 clone SK7 stains a subset of lymphocytes (T cells) and does not stain monocytes and granulocytes (middle plot). Additional analysis of lymphocytes shows that CD3 clone SK7 does not stain any CD19+ B cells (right plot). Details: Normal human whole blood was surface stained with CD3 (clone SK7) and CD19 (clone HIB19). After staining, red blood cells were lysed using 1-step Fix/Lyse Buffer. Cells in the lymphocyte (purple histogram), monocyte (orange histogram), or granulocyte (blue histogram) gates were used to compare CD3 staining (middle plot). Cells in the lymphocyte gate were used to compare CD3 and CD19 staining (right plot). {RE}



CD3 Antibody (35-0036-42) in Flow

Staining of normal human peripheral blood cells with Anti-Human CD19 APC (Product # 17-0199-42) and Mouse IgG1 K Isotype Control PE-Cyanine5-5 (Product # 35-4714-82) (left) or Anti-Human CD3 PE-Cyanine5-5 (right). Cells in the lymphocyte gate were used for analysis.



View more figures on thermofisher.com

Immunohistochemistry (Frozen) (1)

<p>The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society</p> <p>A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates.</p> <p>Authors: Kap YS,van Meurs M,van Driel N,Koopman G,Melief MJ,Brok HP,Laman JD,'t Hart BA</p>	<p>Year 2009</p>
---	-----------------------------

Immunocytochemistry (1)

<p>Clinical cancer research : an official journal of the American Association for Cancer Research</p> <p>Prolongation of overall survival in advanced lung adenocarcinoma patients with the XAGE1 (GAGED2a) antibody.</p> <p>"35-0036 was used in Immunocytochemistry to determine that production of a XAGE1 antibody predicts a good prognosis for patients with lung adenocarcinoma as an immune biomarker."</p> <p>Authors: Ohue Y,Kurose K,Mizote Y,Matsumoto H,Nishio Y,Isobe M,Fukuda M,Uenaka A,Oka M,Nakayama E</p>	<p>Year 2014</p> <p>Species Human</p>
---	---

Flow Cytometry (15)

<p>Frontiers in pharmacology</p> <p>Disruption of PD-1 Enhanced the Anti-tumor Activity of Chimeric Antigen Receptor T Cells Against Hepatocellular Carcinoma.</p> <p>"Published figure using CD3 monoclonal antibody (Product # 35-0036-42) in Flow Cytometry"</p> <p>Authors: Guo X,Jiang H,Shi B,Zhou M,Zhang H,Shi Z,Du G,Luo H,Wu X,Wang Y,Sun R,Li Z</p>	<p>Year 2023</p>
<p>The Journal of biological chemistry</p> <p>Ephrin receptor A10 monoclonal antibodies and the derived chimeric antigen receptor T cells exert an antitumor response in mouse models of triple-negative breast cancer.</p> <p>"Published figure using CD3 monoclonal antibody (Product # 35-0036-42) in Flow Cytometry"</p> <p>Authors: Cha JH,Chan LC,Wang YN,Chu YY,Wang CH,Lee HH,Xia W,Shyu WC,Liu SP,Yao J,Chang CW,Cheng FR,Liu J,Lim SO,Hsu JL,Yang WH,Hortobagyi GN,Lin C,Yang L,Yu D,Jeng LB,Hung MC</p>	<p>Year 2022</p>

View more Flow references on thermofisher.com

More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (i) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (ii) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (iii) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (iv) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.