

# KLRG1 Monoclonal Antibody (13F12F2), Alexa Fluor™ 488, eBioscience™

| Product Details             |   |
|-----------------------------|---|
| Size                        | 100 Tests   |
| Species Reactivity          | Human   |
| Published Species           | Human   |
| Host/Isotype                | Mouse / IgG2a, kappa  |
| Recommended Isotype Control | Mouse IgG2a kappa Isotype Control (eBM2a), Alexa Fluor™ 488, eBioscience™ |
| Class                       | Monoclonal  |
| Type                        | Antibody  |
| Clone                       | 13F12F2   |
| Conjugate                   | Alexa Fluor™ 488  |
| Form                        | Liquid  |
| Concentration               | 5 µL/Test   |
| Purification                | Affinity chromatography   |
| Storage buffer              | PBS, pH 7.2, with 0.1% gelatin, 0.2% BSA                                  |
| Contains                    | 0.09% sodium azide  |
| Storage conditions          | 4° C, store in dark, DO NOT FREEZE!                                       |
| RRID                        | AB_2574451  |

| Applications          | Tested Dilution    | Publications   |
|-----------------------|--------------------|----------------|
| Flow Cytometry (Flow) | 5 µL (0.5 µg)/test | 2 Publications |

## Product Specific Information

Description: This 13F12F2 monoclonal antibody reacts with human killer cell lectin-like receptor subfamily G, member 1 (KLRG1), a type II transmembrane inhibitor receptor of the C-type lectin superfamily that contains an ITIM domain. This inhibitory receptor is expressed on subsets of gamma-delta T cells, NK (CD56dim), CD8+ and CD4+ T Cells. KLRG1 is expressed primarily by cells with an effector/memory phenotype that are short-lived, but capable of immediate effector cell function. Cadherin-E, -N, and -R are ligands for KLRG1. Cadherin/KLRG1 interaction inhibits cytolytic activity and proliferation. The percentage of KLRG1 positive cells can vary considerably, depending on antigen experience.

The clones 13F12F2 and 13A2 appear to recognize a similar epitope based on cross-blocking studies.


Applications Reported: This 13F12F2 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This 13F12F2 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 µL (0.5 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test.

Excitation: 488 nm; Emission: 519 nm; Laser: Blue Laser.

Filtration: 0.2 µm post-manufacturing filtered.

## Product Images For KLRG1 Monoclonal Antibody (13F12F2), Alexa Fluor™ 488, eBioscience™



### KLRG1 Antibody (53-9488-42) in Flow

LEFT: Staining of normal human peripheral blood cells with Anti-Human CD28 eFluor® 450 (Product # 48-0289-42) and Anti-Human KLRG1 Alexa Fluor® 488. CD8+ cells in the lymphocyte gate were used for analysis. RIGHT: Staining of normal human peripheral blood cells with Anti-Human CD56 (NCAM) PE-eFluor® 610 (Product # 61-0567-42) and Anti-Human KLRG1 Alexa Fluor® 488. Cells in the lymphocyte gate were used for analysis.

Flow Cytometry (2)

Nature communications

**IL-1, IL-23, and TGF- drive plasticity of human ILC2s towards IL-17-producing ILCs in nasal inflammation.**

"Published figure using KLRG1 monoclonal antibody (Product # 53-9488-42) in Flow Cytometry"

Authors: Golebski K,Ros XR,Nagasawa M,van Tol S,Heesters BA,Aglmous H,Kradolfer CMA,Shikhagaie MM,Seys S, Hellings PW,van Drunen CM,Fokkens WJ,Spits H,Bal SM

**Species**  
Not Applicable

**Dilution**  
Not Cited

**Year**  
2019

European journal of immunology

**Guidelines for the use of flow cytometry and cell sorting in immunological studies.**

"53-9488 was used in Flow cytometry/Cell sorting to collect the currently accepted best methods for monitoring most of the variation of the major players of immune system."

Authors: Cossarizza A,Chang HD,Radbruch A,Akdis M,Andrà I,Annunziato F,Bacher P,Barnaba V,Battistini L,Bauer WM,Baumgart S,Becher B,Beisker W,Berek C,Blanco A,Borsellino G,Boulais PE,Brinkman RR,Büscher M,Busch DH, Bushnell TP,Cao X,Cavani A,Chattopadhyay PK,Cheng Q,Chow S,Clerici M,Cooke A,Cosma A,Cosmi L,Cumano A, Dang VD,Davies D,De Biasi S,Del Zotto G,Della Bella S,Dellabona P,Deniz G,Dessing M,Diefenbach A,Di Santo J,Dieli F,Dolf A,Donnenberg VS,Dörner T,Ehrhardt GRA,Endl E,Engel P,Engelhardt B,Esser C,Everts B,Dreher A,Falk CS, Fehniger TA,Filby A,Fillatreau S,Follo M,Förster I,Foster J,Foulds GA,Frenette PS,Galbraith D,Garbi N,García-Godoy MD,Geginat J,Ghorechi K,Gibellini L,Goettlinger C,Goodyear CS,Gori A,Grogan J,Gross M,Grützkau A,Grummitt D, Hahn J,Hammer Q,Hauser AE,Haviland DL,Hedley D,Herrera G,Herrmann M,Hiepe F,Holland T,Hombrink P,Houston JP,Hoyer BF,Huang B,Hunter CA,Iannone A,Jäck HM,Jávega B,Jonjic S,Juelke K,Jung S,Kaiser T,Kalina T,Keller B, Khan S,Kienhöfer D,Kroneis T,Kunkel D,Kurts C,Kvistborg P,Lannigan J,Lantz O,Larbi A,LeibundGut-Landmann S, Leipold MD,Levings MK,Litwin V,Liu Y,Lohoff M,Lombardi G,Lopez L,Lovett-Racke A,Lubbbers E,Ludewig B,Lugli E, Maecker HT,Martrus G,Matarese G,Maueröder C,McGrath M,McInnes I,Mei HE,Melchers F,Melzer S,Mielenz D,Mills K, Mirrer D,Mjösberg J,Moore J,Moran B,Moretta A,Moretta L,Mosmann TR,Müller S,Müller W,Münz C,Multhoff G,Munoz LE,Murphy KM,Nakayama T,Nasi M,Neudörfl C,Nolan J,Nourshargh S,O'Connor JE,Ouyang W,Oxenius A,Palankar R, Panse I,Peterson P,Peth C,Petritz J,Philips D,Pickl W,Piconese S,Pinti M,Pockley AG,Podolska MJ,Pucillo C,Quataert SA,Radstake TRDJ,Rajwa B,Rebhahn JA,Recktenwald D,Remmerswaal EBM,Rezvani K,Rico LG,Robinson JP, Romagnani C,Rubartelli A,Ruckert B,Ruland J,Sakaguchi S,Sala-de-Oyanguren F,Samstag Y,Sanderson S,Sawitzki B, Scheffold A,Schiemann M,Schildberg F,Schimisky E,Schmid SA,Schmitt S,Schober K,Schüler T,Schulz AR, Schumacher T,Scotta C,Shankey TV,Shemer A,Simon AK,Spidlen J,Stall AM,Stark R,Stehle C,Stein M,Steinmetz T, Stockinger H,Takahama Y,Tarnok A,Tian Z,Toldi G,Tornack J,Traggiai E,Trotter J,Ulrich H,van der Braber M,van Lier RAW,Veldhoen M,Vento-Asturias S,Vieira P,Voehringer D,Volk HD,von Volkman K,Waisman A,Walker R,Ward MD, Warnatz K,Warth S,Watson JV,Watzl C,Wegener L,Wiedemann A,Wienands J,Willimsky G,Wing J,Wurst P,Yu L,Yue A, Zhang Q,Zhao Y,Ziegler S,Zimmermann J

**Species**  
Human

**Dilution**  
Not Cited

**Year**  
2017

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.