



# CD107a (LAMP-1) Monoclonal Antibody (eBioH4A3), PE-eFluor™ 610, eBioscience™

<b>Product Details</b>	
Size	100 Tests
Species Reactivity	Human
Published Species	Rhesus monkey
Host/Isotype	Mouse / IgG1, kappa
Recommended Isotype Control	Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), PE-eFluor™ 610, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	eBioH4A3
Conjugate	PE-eFluor™ 610
Excitation/Emission Max	565/606 nm
Form	Liquid
Concentration	5 μL/Test
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.2% BSA
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2574572

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	5 μL (0.25 μg)/test	9 Publications

#### **Product Specific Information**

Description: The eBioH4A3 monoclonal antibody reacts with human CD107a, also known as lysosomal-associated membrane protein-1 (LAMP-1). CD107a is a highly glycosylated protein of approximately 110kDa. It is predominantly expressed intracellularly in the lysosomal/endosomal membrane in nearly all cells. CD107a is transiently expressed on the cell surface of degranulating cytolytic T cells, and is also upregulated on the surface of activated platelets and some cancer cells.

Applications Reported: This eBioH4A3 antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

Applications Tested: This eBioH4A3 antibody has been pre-titrated and tested by intracellular staining and flow cytometric analysis of Jurkat cells using the Intracellular Fixation & Permeabilization Buffer Set (cat. 88-8824) and protocol. Please refer to Best Protocols: Protocol A: Two step protocol for (cytoplasmic) intracellular proteins located under the Resources Tab online. This can be used at 5 µL (0.25 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 μL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test.

PE-eFluor® 610 can be excited with laser lines from 488-561 nm and emits at 607 nm. We recommend using a 610/20 band pass filter (equivalent to PE-Texas Red®). Please make sure that your instrument is capable of detecting this fluorochome.


Light sensitivity: This tandem dye is sensitive to photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100  $\mu$ L of cell sample + 100  $\mu$ L of IC Fixation Buffer) or 1-step Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency /compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

Excitation: 488-561 nm; Emission: 607 nm; Laser: Blue Laser, Green Laser, Yellow-Green Laser.

Filtration: 0.2 µm post-manufacturing filtered.

# Product Images For CD107a (LAMP-1) Monoclonal Antibody (eBioH4A3), PE-eFluor™ 610, eBioscience™



# CD107a (LAMP-1) Antibody (61-1079-42) in Flow

Intracellular staining of Jurkat cells with Mouse IgG1 K Isotype Control PE-eFluor® 610 (Product # 61-4714) (blue histogram) or Anti-Human CD107a (LAMP-1) PE-eFluor® 610 (purple histogram) using the Intracellular Fixation & Permeabilization Buffer Set (Product # 88-8824) and protocol. Total viable cells, as determined by Fixable Viability Dye eFluor® 660 (Product # 65-0864), were used for analysis.

View more figures on thermofisher.com

#### □ 9 References

# Flow Cytometry (9)

Nature microbiology

# CD8<sup>+</sup> lymphocytes do not impact SIV reservoir establishment under ART.

"61-1079-42 was used in Flow cytometry/Cell sorting to indicate that during early SIV infection, the viral reservoir that persists under ART is established largely independent of CTL control."

Authors: Statzu M,Jin W,Fray EJ,Wong AKH,Kumar MR,Ferrer E,Docken SS,Pinkevych M,McBrien JB,Fennessey CM, Keele BF,Liang S,Harper JL,Mutascio S,Franchitti L,Wang H,Cicetti D,Bosinger SE,Carnathan DG,Vanderford TH, Margolis DM,Garcia-Martinez JV,Chahroudi A,Paiardini M,Siliciano J,Davenport MP,Kulpa DA,Siliciano RS,Silvestri G

**Year** 2023

Species Rhesus monkey

Signal transduction and targeted therapy

# Interaction between HLA-G and NK cell receptor KIR2DL4 orchestrates HER2-positive breast cancer resistance to trastuzumab.

"Published figure using CD107a (LAMP-1) monoclonal antibody (Product # 61-1079-42) in Flow Cytometry" Authors: Zheng G,Guo Z,Li W,Xi W,Zuo B,Zhang R,Wen W,Yang AG,Jia L

**Year** 2021

View more Flow references on thermofisher.com

### More applications with references on thermofisher.com