CD4 Monoclonal Antibody (OKT4 (OKT-4)), Super Bright[™] 436, eBioscience[™]

| Product Details | |
|--------------------------------|---|
| Size | 100 Tests |
| Species Reactivity | Human |
| Host/Isotype | Mouse / IgG2b, kappa |
| Recommended Isotype Control | Mouse IgG2b kappa Isotype Control (eBMG2b), Super Bright™ 436, eBioscience™ |
| Class | Monoclonal |
| Туре | Antibody |
| Clone | OKT4 (OKT-4) |
| Conjugate | Super Bright™ 436 |
| Excitation/Emission Max | 413/431 nm |
| Form | Liquid |
| Concentration | 5 µL/Test |
| Purification | Affinity chromatography |
| Storage buffer | PBS, pH 7.2, with BSA |
| Contains | 0.09% sodium azide |
| Storage conditions | 4° C, store in dark, DO NOT FREEZE! |
| RRID | AB_2744764 |
| | |

| Applications | Tested Dilution | Publications |
|-----------------------|----------------------|----------------|
| Flow Cytometry (Flow) | 5 μL (0.125 μg)/test | 7 Publications |

Product Specific Information

Description: The OKT4 monoclonal antibody reacts with human CD4, a 59 kDa cell surface glycoprotein expressed by the majority of thymocytes, a subpopulation of mature T cells (T-helper cells) and in low levels on monocytes. CD4 is a receptor for the human immunodeficiency virus (HIV). The OKT4 antibody recognizes a different epitope than the RPA-T4 monoclonal antibody, and these antibodies do not cross-block binding to each other's respective epitopes.

Applications Reported: This OKT4 (OKT-4) antibody has been reported for use in flow cytometric analysis.

Applications Tested: This OKT4 (OKT-4) antibody has been pre-titrated and tested by flow cytometric analysis of normal human blood peripheral blood cells. This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

Super Bright 436 can be excited with the violet laser line (405 nm) and emits at 436 nm. We recommend using a 450/50 bandpass filter, or equivalent. Please make sure that your instrument is capable of detecting this fluorochrome.


When using two or more Super Bright dye-conjugated antibodies in a staining panel, it is recommended to use Super Bright Complete Staining Buffer (Product # SB-4401) to minimize any non-specific polymer interactions. Please refer to the datasheet for Super Bright Staining Buffer for more information.

1

Excitation: 405 nm; Emission: 436 nm; Laser: Violet Laser

Super Bright Polymer Dyes are sold under license from Becton, Dickinson and Company.

Product Images For CD4 Monoclonal Antibody (OKT4 (OKT-4)), Super Bright[™] 436, eBioscience[™]



CD4 Antibody (62-0048-42) in Flow

Normal human peripheral blood cells were stained with CD19 Monoclonal Antibody, FITC (Product # 11-0119-82) and Mouse IgG2b kappa Isotype Control, Super Bright 436 (Product # 62-4724-82) (left) or CD4 Monoclonal Antibody, Super Bright 436 (right). Cells in the lymphocyte gate were used for analysis.

View more figures on thermofisher.com

7 References

Flow Cytometry (7)

| Disease markers Estrogen Protects against Renal Ischemia-Reperfusion Injury by Regulating Th17/Treg Cell Immune Balance. | Year 2022 |
|--|---------------------|
| "Published figure using CD4 monoclonal antibody (Product # 62-0048-42) in Flow Cytometry" Authors: Zhang Y,Chang Y,Han Z,Ma K,Zeng X,Li L | |
| Pathogens & immunity | Year |
| | 2022 |
| Plasma Extracellular Vesicle Subtypes May be Useful as Potential | |
| Biomarkers of Immune Activation in People With HIV. | |
| "Published figure using CD4 monoclonal antibody (Product # 62-0048-42) in Flow Cytometry" | |

Authors: Bazié WW,Boucher J,Vitry J,Goyer B,Routy JP,Tremblay C,Trottier S,Jenabian MA,Provost P,Alary M,Gilbert C

View more Flow references on thermofisher.com

More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is more of the product set of any product will conform to such model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No OTHER WARRANTES, EXPRESS OR INVERDED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTES OF MERCHANTABILITY, FITNESS FOR ANY PARTICURST OR NON INFRINGEMENT. BUYER'S SCILUSVIE CEMELOY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTE, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS SOLEN OF THOUR CEMANTER IN CONFORMING PRODUCTS SOLEN OF THOUR CESS OLE OFTION. THERE IS NO BILICATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS SOLEN OF THOUR CESS OLE OFTION. THERE IS NO BILICATION TO FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commencial uses, in vitro diagnostic uses, arvivo or in vivo therapeutic uses, orany type of consumption to human or animals.