

IL-8 (CXCL8) Monoclonal Antibody (3IL8-H10)

Product Details	
Size	500 µg
Species	Human
Published Species	Human
Expression System	Mouse / IgG1
Class	Monoclonal
Type	Antibody
Clone	3IL8-H10
Conjugate	Unconjugated
Immunogen	Recombinant human IL-8
Form	Liquid
Concentration	1.0 mg/mL
Purification	Protein G
Storage buffer	PBS
Contains	no preservative
Storage Conditions	-20°C
RRID	AB_223583

Applications	Tested Dilution	Publications
ELISA (ELISA)	Assay Dependent	18 Publications
Immunocytochemistry (ICC)	2 µg/mL	-
Immunofluorescence (IF)	2 µg/mL	-
Western Blot (WB)	Assay Dependent	-
Immunohistochemistry (IHC)	-	1 Publication
Radioimmune Assays (RIA)	-	1 Publication

Product Specific Information

M801 targets IL-8 in ELISA, and WB applications and shows reactivity with Human samples.

The M801 immunogen is recombinant human IL-8.


M801 detects IL-8 which has a predicted molecular weight of approximately 9 kDa.

The M801 IL8 antibody (clone 3IL8-H10) has successfully been paired as the coating antibody in a sandwich ELISA with detection antibody M802B (biotinylated conjugate of clone I8-S2). Typical dilutions for sandwich ELISA include 1 µg/mL for coating and 0.125 - 0.25 µg/mL for detection.

Antibody M801 (clone 3IL8-H10) and biotinylated antibody M802B (clone I8-S2) have successfully been used in combination with recombinant IL8 protein SIL8 in ELISA applications.

This product has been tested for endotoxins by limulus amoebocyte lysate (LAL) assay and contains an endotoxin concentration of less than or equal to 10 endotoxin units per milligram (EU/mg).


Advanced Verification Data



IL-8 (CXCL8) Antibody (M801)

Modulation of expression of target protein by cell treatment to demonstrate antibody specificity. Immunofluorescence analysis of CXCL8 using anti-CXCL8 mouse monoclonal antibody (Catalog # M801) shows increased expression of CXCL8 in U87MG upon treatment with thapsigargin. Cell treatment validation info.

Product Images For IL-8 (CXCL8) Monoclonal Antibody (3IL8-H10)




IL-8 (CXCL8) Antibody (M801) in IF

Immunofluorescence analysis of IL-8 was performed using 70% confluent log phase U-87 MG cells treated 1 uM thapsigargin for 24 hours. The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 10 minutes, and blocked with 1% BSA for 1 hour at room temperature. The cells were labeled with IL-8 (3IL8-H10) Mouse Monoclonal Antibody (Product # M801) at 2 µg /mL in 0.1% BSA and incubated for 3 hours at room temperature and then labeled with Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate (Product # A28175) at a dilution of 1:2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI (Product # S36938). F-actin (Panel c: red) was stained with Alexa Fluor® 555 Rhodamine Phalloidin (Product # R415, 1:300). Panel d represents the merged image showing cytoplasmic localization. Panel e is untreated cell with no signal. Panel f represents control cells with no primary antibody to assess background. The images were captured at 60X magnification.

IL-8 (CXCL8) Antibody (M801) in ELISA

Sandwich ELISA analysis of human IL-8 was performed using a Human IL-8 Colorimetric ELISA kit (Product # EH2IL8) by loading 50 µL per well of Human IL-8 Recombinant Protein (Product # SIL8) in dodecuplicate at 1000, 400, 160, 64, 25.6, and 0 pg/mL across a 3 µg/mL mouse anti-Human IL-8 (Product # M801) pre-coated plate and incubating for 1 hour at room temperature. The plate was washed, and then incubated with 50 µL per well of a biotinylated mouse anti-human IL-8 monoclonal antibody (Product # M802B) in duplicate at 0.25 µg/mL for 1 hour at room temperature. The plate was washed and incubated with 100 µL per well of Streptavidin-HRP (Product # N504) in all test wells at a 1:4,000 dilution for 30 minutes at room temperature. Detection was performed using 1-Step Ultra TMB substrate (Product # 34028) for 30 minutes at room temperature in the dark. The plate was then stopped with 0.16M sulfuric acid. Absorbances were read on a spectrophotometer at 450-550 nm.



View more figures on [thermofisher.com](https://www.thermofisher.com)

ELISA (18)

Journal of immunology research

The Antiseptic Octenidine Inhibits Langerhans Cell Activation and Modulates Cytokine Expression upon Superficial Wounding with Tape Stripping.

"M801 was used in an ELISA assay to investigate the influence of octenidine on epidermal Langerhans cells and cytokine secretion pattern of skin cells during wound healing in a model without disruption of the normal skin structure."

Authors: Nikoli N,Kienzl P,Tajpara P,Vierhapper M,Matiasek J,Elbe-Bürger A

Species
Human

Dilution
Not Cited

Year
2019

FASEB journal : official publication of the Federation of American Societies for Experimental Biology

Resolvin D2 decreases TLR4 expression to mediate resolution in human monocytes.

"Published figure using IL-8 (CXCL8) monoclonal antibody (Product # M801) in ELISA"

Authors: Croasdell A,Sime PJ,Phipps RP

Species
Not Applicable

Dilution
Not Cited

Year
2016

[View more ELISA references on thermofisher.com](#)

Radioimmune Assays (1)

Radiation research

The human salivary proteome is radiation responsive.

"Published figure using IL-8 (CXCL8) monoclonal antibody (Product # M801) in Radioimmune Assays"

Authors: Moore HD,Ivey RG,Voytovich UJ,Lin C,Stirewalt DL,Pogosova-Agadjanyan EL,Paulovich AG

Species
Not Applicable

Dilution
Not Cited

Year
2014

Immunohistochemistry (1)

Journal of immunology (Baltimore, Md. : 1950)

Rapamycin treatment depresses intragraft expression of KC/MIP-2, granzyme B, and IFN-gamma in rat recipients of cardiac allografts.

"M801 was used in immunohistochemistry to investigate the effects of rapamycin on KC/MIP-2, granzyme B, and interferon gamma expression on cardiac allografts"

Authors: Wieder KJ,Hancock WW,Schmidbauer G,Corpier CL,Wieder I,Kobzik L,Strom TB,Kupiec-Weglinski JW

Species
Human

Dilution
Not Cited

Year
1993

More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.