



hnRNP K Recombinant Rabbit Monoclonal Antibody (ARC0512)


Product Details	
Size	100 μL
Species Reactivity	Human, Mouse, Rat
Host/Isotype	Rabbit / IgG
Expression system	HEK293 cells
Class	Recombinant Monoclonal
Туре	Antibody
Clone	ARC0512
Conjugate	Unconjugated
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human hnRNP K (P61978)
Form	Liquid
Concentration	0.6 mg/mL
Purification	Affinity Chromatography
Storage buffer	PBS, pH 7.3, with 50% glycerol, 0.05% BSA
Contains	0.02% sodium azide
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles
RRID	AB_2848963

Applications	Tested Dilution	Publications
Western Blot (WB)	1:500-1:2,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:500-1:1000	-
Immunocytochemistry (ICC/IF)	1:50-1:200	-

Product Specific Information


Immunogen sequence: METEQPEETF PNTETNGEFG KRPAEDMEEE QAFKRSRNTD EMVELRILLQ SKNAGAVIGK GGKNIKALRT DYNASVSVPD SSGPERILSI SADIETIGEI

Product Images For hnRNP K Recombinant Rabbit Monoclonal Antibody (ARC0512)




hnRNP K Antibody (MA5-35058) in WB

Western blot analysis of hnRNP K in extracts of various cell lines. Samples were incubated with hnRNP K Monoclonal antibody (Product # MA5-35058) using a dilution of 1:2,000, followed by HRP Goat Anti-Rabbit IgG (H+L) at a dilution of 1: 10,000. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.



hnRNP K Antibody (MA5-35058) in ICC/IF

Confocal imaging (Immunocytochemistry) of hnRNP K in U-2 OS cells. Samples were incubated with hnRNP K Monoclonal antibody (Product # MA5-35058) using a dilution of 1:100 (Red). The cells were counterstained with alpha Tubulin Mouse mAb (dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 60x.



hnRNP K Antibody (MA5-35058) in IHC (P)

Immunohistochemistry analysis of hnRNP K in paraffin-embedded human esophageal. Samples were incubated with hnRNP K Monoclonal antibody (Product # MA5-35058) using a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

View more figures on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPERESS OR IMPLEA REPRESS OR IMPLEA REGRANTED INITY, IMPLIED WARRANTIES, EXPERESS OR IMPLEA REPRESS OR IMPLEAD IN INFINITY ("INFINITY PROPOSE"). AND ANY PARTICULAR VARRANTIES, EXPERESS OR IMPLEAD IN INFINITY PROPOSED, OR NON INFINITY PROPOSED IN INFINITY PROPOSED I