

mTOR Polyclonal Antibody

Product Details	
Size	100 µg
Species Reactivity	Human, Mouse, Non-human primate, Rat
Published Species	Rat, Human
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Type	Antibody
Conjugate	Unconjugated
Immunogen	Immunogen corresponds to amino acids 1223-1290 of human mTOR.
Form	Liquid
Concentration	1 mg/mL
Purification	Protein G
Storage buffer	PBS with 1mg/mL BSA, 30% glycerol
Contains	0.05% sodium azide
Storage conditions	-20°C
RRID	AB_325371


Applications	Tested Dilution	Publications
Western Blot (WB)	1:100-1:1,000	5 Publications
Immunocytochemistry (ICC/IF)	1:50-1:200	-
Immunoprecipitation (IP)	2 µg	-

Product Specific Information

3-8% Tris-Acetate Gel is recommended for Western blot.

PA1-518 detects a predominant band at ~290kD corresponding to mTOR. Other lower MW nonspecific bands of unknown identity were also detected at ~65kD and ~35kD in some cell lysates.

Product Images For mTOR Polyclonal Antibody




mTOR Antibody (PA1-518)

IP-MS enrichment of MTOR (LFQ intensity): MTOR was enriched 74-fold from HCT116 lysate compared to background proteins, using the optimized IP-MS workflow with Pierce MS-Compatible Magnetic IP Kit protein A/G (Product # 90409) and MTOR antibody (Product # PA1-518). See more information on IP-MS verification of antibody selectivity. {IP-MS}


mTOR Antibody (PA1-518) in WB

Western blot was performed using Anti-mTOR Polyclonal Antibody (Product # PA1-518) and a ~280kDa band corresponding to Serine/threonine-protein kinase mTOR was observed across cell lines and tissues tested . Whole cell extracts (30 µg lysate) of SiHa (Lane 1), K-562 (Lane 2), HEK-293 (Lane 3), MCF7 (Lane 4), MDA-MB-231 (Lane 5), HEL 92.1.7 (Lane 6), Hep G2 (Lane 7), Mouse Stomach (Lane 8), Rat Cerebellum (Lane 9), Mouse Heart (Lane 10) were electrophoresed using NuPAGE™ 3-8% Tris-Acetate Protein Gel (Product # EA0378BOX). Resolved proteins were then transferred onto a Nitrocellulose membrane (Product # LC2002) by iBlot® 2 Dry Blotting System (Product # IB21001). The blot was probed with the primary antibody (1:1000) and detected by chemiluminescence with Goat anti-Rabbit IgG (Heavy Chain) Superclonal™ Recombinant Secondary Antibody, HRP (Product # A27036, 1:4000) using the iBright FL 1000 (Product # A32752). Chemiluminescent detection was performed using Novex® ECL Reagent Kit (Product # WP20005).



mTOR Antibody (PA1-518) in ICC/IF

Immunofluorescent analysis of mTOR (green) in HeLa cells. The cells were fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton X-100 in PBS, and blocked with 1% Blocker BSA (Product # 37525), each for 15 minutes at room temperature. Cells were stained with an mTOR polyclonal antibody (Product # PA1-518) at a concentration of 10 µg/mL in 1% Blocker BSA in PBS (right panel), or incubated in blocking buffer alone as a negative control (left panel) overnight at 4C, and then incubated with a Dylight 488-conjugated goat anti-rabbit IgG secondary antibody (Product # 35552) at a dilution of 1:1000 for 1 hour at room temperature. F-Actin (red) was stained with a DyLight 554-conjugated phalloidin control (Product # 21834) and nuclei (blue) were stained with DAPI (Product # 46190).



View more figures on thermofisher.com

Western Blot (5)

<p>Cell death & disease</p> <p>PTPN14 promotes gastric cancer progression by PI3KA/AKT/mTOR pathway.</p> <p>"PA1-518 was used in Western blotting following immunoprecipitation to provide a theoretical basis to better understand the occurrence and development of gastric cancer."</p> <p>Authors: Li H,Guan B,Liu S,Liu H,Song L,Zhang G,Zhao R,Zhou C,Gao P</p>	<p>Year 2023</p> <p>Species Human</p>
<p>Pharmaceuticals (Basel, Switzerland)</p> <p>Partial Synthetic PPARγ Derivative Ameliorates Aorta Injury in Experimental Diabetic Rats Mediated by Activation of miR-126-5p Pi3k /AKT/PDK 1/mTOR Expression.</p> <p>"PA1-518 was used in Western Blotting to reveal that HFD/STZ-induced T2D increased fasting blood glucose, ET-1, ICAM-1, E-selectin, and VIP levels, while decreasing the expression of both microRNA126-5p and p-AKT/p-Pi3k/p-PDK-1/p-mTOR phosphorylation."</p> <p>Authors: Ahmed YM,Orfali R,Abdelwahab NS,Hassan HM,Rateb ME,AboulMagd AM</p>	<p>Year 2022</p> <p>Species Rat</p>

[View more WB references on thermofisher.com](#)

More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.