




NDUFS2 Polyclonal Antibody

Product Details	
Size	100 μL
Species Reactivity	Cynomolgus monkey, Human
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Conjugate	Unconjugated
Immunogen	E. coli-derived Human NDUFS2 fragment
Form	Liquid
Concentration	1 mg/mL
Storage conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.
RRID	AB_2867014


Applications	Tested Dilution	Publications
Western Blot (WB)	1:500-1:2,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:50-1:200	-
Immunoprecipitation (IP)	1-5 µL/mg lysate	-

Product Images For NDUFS2 Polyclonal Antibody



NDUFS2 Antibody (PA5-112278) in IHC (P)

Immunohistochemistry analysis of NDUFS2 in formalin-fixed paraffin embedded sections cynomolgus heart tissue sections using NDUFS2 Polyclonal Antibody (Product # PA5-112278) at a dilution of 1:100.



NDUFS2 Antibody (PA5-112278) in WB

Western Blot analysis of NDUFS2 using NDUFS2 Polyclonal Antibody (Product # PA5-112278) at a dilution of 1:500. Lane A: HeLa whole cell lysate, Lane B: Jurkat whole cell lysate, Lane C: 293T whole cell lysate, (Lysates/proteins at 30 µg per lane). The secondary antibody used was a Goat Anti-Rabbit IgG (H+L) /HRP at 1:10,000 dilution. Developed using the ECL technique and performed under reducing conditions. Predicted band size: 53 kDa. Observed band size: 50 kDa.



NDUFS2 Antibody (PA5-112278) in IP

Immunoprecipitation of NDUFS2 was performed on (Lane A) 0.5 mg Jurkat whole cell lysate using 4 μL of NDUFS2 Polyclonal Antibody (Product # PA5-112278) at a dilution of 1:100, and 60 μg of Immunomagnetic beads Protein A/G. A Clean-Blot IP Detection Reagent (HRP) was used as a secondary antibody at a dilution of 1:1,000. Developed using the ECL technique and performed under reducing conditions. Predicted band size: 53 kDa. Observed band size: 45 kDa.