




Phospho-NFkB p50/p105 (Ser923) Polyclonal Antibody


Product Details	
Size	100 μL
Species Reactivity	Human, Mouse, Rat
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human NF-kB p105 around the phosphorylation site of Ser923.
Form	Liquid
Concentration	1 mg/mL
Storage conditions	-20°C
RRID	AB_2903105

Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000-1:3,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:50-1:200	-
Immunocytochemistry (ICC/IF)	1:100-1:500	-


Product Images For Phospho-NFkB p50/p105 (Ser923) Polyclonal Antibody



Phospho-NFkB p50/p105 (Ser923) Antibody (PA5-118605) in ICC/IF Immunocytochemical analysis of Phospho-NFkB p50/p105 (Ser923) in Hela cells (treated for 4 hours with LPS) using a Phospho-NFkB p50/p105 (Ser923) polyclonal antibody (Product # PA5-118605). The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with the primary antibody and mouse anti-beta tubulin antibody at a dilution of 1:200 for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody. The nuclear counter stain is DAPI(blue).



Phospho-NFkB p50/p105 (Ser923) Antibody (PA5-118605) in IHC (P) Immunohistochemical analysis of Phospho-NFkB p50/p105 (Ser923) in Human kidney tissue using a Phospho-NFkB p50/p105 (Ser923) polyclonal antibody (Product # PA5-118605). The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.



Phospho-NFkB p50/p105 (Ser923) Antibody (PA5-118605) in IHC (P) Immunohistochemical analysis of Phospho-NFkB p50/p105 (Ser923) in Mouse kidney tissue using a Phospho-NFkB p50/p105 (Ser923) polyclonal antibody (Product # PA5-118605). The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.

View more figures on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED. A FERDIO IS LI