



# **Vimentin Polyclonal Antibody**


<b>Product Details</b>		
Size	100 μL	
Species Reactivity	Human, Mouse	
Host/Isotype	Rabbit / IgG	
Class	Polyclonal	
Туре	Antibody	
Conjugate	Unconjugated	
Immunogen	Recombinant protein corresponding to Human VIM	
Form	Liquid	
Concentration	0.2 mg/mL	
Purification	Anitgen affinity chromatography, Antigen affinity chromatography	
Storage buffer	PBS, pH 7.2, with 40% glycerol	
Contains	0.02% sodium azide	
Storage conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.	
RRID	AB_2789129	

Applications	Tested Dilution	Publications
Western Blot (WB)	0.04-0.4 μg/mL	-
Immunohistochemistry (IHC)	1:1,000-1:2,500	-
Immunocytochemistry (ICC/IF)	0.25-2 μg/mL	-

### **Product Specific Information**


Immunogen sequence: FANYIDKVRF LEQQNKILLA ELEQLKGQGK SRLGDLYEEE MRELRRQVDQ LTNDKARVEV ERDNLAEDIM RLREKLQEEM LQREEAENTL QSFRQDVDNA SLARLDLERK VES

#### **Product Images For Vimentin Polyclonal Antibody**



#### Vimentin Antibody (PA5-81968)

Relative expression in western blot: Antibody specificity was demonstrated by detection of known differential basal expression of the target across tissue/cell models. Expression of Vimentin was observed specifically in U-251MG cells and in MCF-7 cells using anti-Vimentin Polyclonal Antibody (Product #PA5-81968) in western blot. The relative expression levels of Vimentin within each cell line is shown using RNA-Seq. {RE}



## Vimentin Antibody (PA5-81968)

Relative expression in different tissues in IHC: Detection of differential expression levels of Vimentin demonstrates antibody specificity. Immunohistochemical analysis of Vimentin using anti-Vimentin Polyclonal Antibody (Product #PA5-81968), shows significant staining of Vimentin in ovary and shows minimal or weak staining in skeletal muscle tissues. The relative expression levels of Vimentin within each tissue is shown using RNA-Seq. {RE}

RNA [TPM]

800 1600 2400 3200 4000



For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless of services the use of subject to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CORPORMING PRODUCTS DURING THE WARRANTY PERFIOI DIS LIMITED. A REFLION FOR THE NON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying is.