



UBE1 Polyclonal Antibody


Product Details		
Size	100 μL	
Species Reactivity	Human, Mouse, Rat	
Host/Isotype	Rabbit / IgG	
Class	Polyclonal	
Type	Antibody	
Conjugate	Unconjugated	
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 300-600 of human UBA1 (NP_6950121)	
Form	liquid	
Concentration	0.57 mg/mL	
Purification	Affinity Chromatography	
Storage buffer	PBS, pH 7.3, with 50% glycerol	
Contains	0.01% thimerosal	
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles	
RRID	AB_2804952	

Applications	Tested Dilution	Publications
Western Blot (WB)	1:500-1:2,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:50-1:200	-
Immunocytochemistry (ICC/IF)	1:50-1:200	-

Product Specific Information


Immunogen sequence: KISFKSLVAS LAEPDFVVTD FAKFSRPAQL HIGFQALHQF CAQHGRPPRP RNEEDAAELV ALAQAVNARA LPAVQQNNLD EDLIRKLAYV AAGDLAPINA FIGGLAAQEV MKACSGKFMP IMQWLYFDAL ECLPEDKEVL TEDKCLQRQN RYDGQVAVFG SDLQEKLGKQ KYFLVGAGAI GCELLKNFAM IGLGCGEGGE IIVTDMDTIE KSNLNRQFLF RPWDVTKLKS DTAAAAVRQM NPHIRVTSHQ NRVGPDTERI YDDDFFQNLD GVANALDNVD ARMYMDRRCV YYRKPLLESG T; Positive Samples: U-251MG, Rat spleen; Cellular Location: Cytoplasm, Mitochondrion, Nucleus, Nucleus

Product Images For UBE1 Polyclonal Antibody




UBE1 Antibody (PA5-88493) in WB

Western blot analysis of UBE1 in extracts of various cell lines. Samples were incubated with UBE1 Polyclonal antibody (Product # PA5-88493) using a dilution of 1:3,000, followed by HRP Goat Anti-Rabbit IgG (H+L) at a dilution of 1:10,000. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.



UBE1 Antibody (PA5-88493) in ICC/IF

Immunofluorescence analysis of UBE1 in HeLa cells. Samples were incubated with UBE1 Polyclonal antibody (Product # PA5-88493) using a dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



UBE1 Antibody (PA5-88493) in IHC (P)

Immunohistochemistry analysis of UBE1 in paraffin-embedded human colon carcinoma. Samples were incubated with UBE1 Polyclonal antibody (Product # PA5-88493) using a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

View more figures on thermofisher.com

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED AND THE PRIOD CONFORMING PRODUCTS AND THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS; AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS; AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS AND THE PRODUCTS