




TGM2 Polyclonal Antibody

Product Details	
Size	100 μg
Species Reactivity	Human
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Conjugate	Unconjugated
Immunogen	Recombinant Human Protein-glutamine gamma-glutamyltransferase 2 protein (4-544AA)
Form	Liquid
Concentration	3 mg/mL
Purification	Protein G
Storage buffer	PBS, pH 7.4, with 50% glycerol
Contains	0.03% ProClin 300
Storage conditions	-20°C or -80°C if preferred
RRID	AB_2812921


Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000-1:5,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:200-1:500	-
Immunocytochemistry (ICC/IF)	1:50-1:200	-
ELISA (ELISA)	Assay-Dependent	-

Product Images For TGM2 Polyclonal Antibody




TGM2 Antibody (PA5-98308) in ICC/IF

Immunofluorescent analysis of TGM2 in Hela cells using a TGM2 polyclonal antibody (Product # PA5-98308) at a dilution of 1:133. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated Goat Anti-Rabbit IgG(H+L). Cells were counter-stained with DAPI.



TGM2 Antibody (PA5-98308) in IHC (P)

Immunohistochemical analysis of TGM2 in paraffin embedded human liver cancer using a TGM2 polyclonal antibody (Product # PA5-98308) at a dilution of 1:400. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



TGM2 Antibody (PA5-98308) in WB

Western Blot analysis of TGM2 using a TGM2 Polyclonal antibody (Product # PA5-98308) at a concentration of 4 μ g/mL. Positive WB detected in: Hela whole cell lysate, A549 whole cell lysate, HepG2 whole cell lysate. A secondary Goat polyclonal antibody to rabbit IgG was applied at a 1:50,000 dilution. Observed band size: 78 kDa.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect, as set forth in the Production documentation, specifications and/or accompanying package intentation?). No claim of suitability for use in applications, the warranty provided herein is valid only when used by properly trained individuals. Intoins otherwise stated in the Decumentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTENBES FOR ANY PARTICULAR PURPOSE, OR NON INFIRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S), AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S), AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACEMENT OF OR REFUND FOR THE RODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation to human or animals.