




MOV10L1 Polyclonal Antibody

| Product Details | | |
|------------------------|--|--|
| Size | 100 μg | |
| Species Reactivity | Human | |
| Host/Isotype | Rabbit / IgG | |
| Class | Polyclonal | |
| Туре | Antibody | |
| Conjugate | Unconjugated | |
| Immunogen | Recombinant Human RNA helicase Mov10l1 protein (336-425AA) | |
| Form | Liquid | |
| Concentration | 4.06 mg/mL | |
| Purification | Protein G | |
| Storage buffer | PBS, pH 7.4, with 50% glycerol | |
| Contains | 0.03% ProClin 300 | |
| Storage conditions | -20°C or -80°C if preferred | |
| RRID | AB_2813511 | |


| Applications | Tested Dilution | Publications |
|---|-----------------|--------------|
| Western Blot (WB) | 1:500-1:5,000 | - |
| Immunohistochemistry (Paraffin) (IHC (P)) | 1:200-1:500 | - |
| Immunocytochemistry (ICC/IF) | 1:50-1:200 | - |
| ELISA (ELISA) | Assay-Dependent | - |

Product Images For MOV10L1 Polyclonal Antibody




MOV10L1 Antibody (PA5-98898) in WB

Western Blot analysis of MOV10L1 using a MOV10L1 Polyclonal antibody (Product # PA5-98898) at a concentration of 6.5 μ g/mL. Positive WB detected in: Hela whole cell lysate, PC-3 whole cell lysate. A secondary Goat polyclonal antibody to rabbit IgG was applied at a 1:50,000 dilution. Observed band size: 136 kDa.



MOV10L1 Antibody (PA5-98898) in ICC/IF

Immunofluorescent analysis of MOV10L1 in HepG2 cells using a MOV10L1 polyclonal antibody (Product # PA5-98898) at a dilution of 1:133. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated Goat Anti-Rabbit IgG(H+L). Cells were counter-stained with DAPI.



MOV10L1 Antibody (PA5-98898) in IHC (P)

Immunohistochemical analysis of MOV10L1 in paraffin embedded human breast cancer using a MOV10L1 polyclonal antibody (Product # PA5-98898) at a dilution of 1:400. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

View more figures on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPERESS OR IMPLEA REPRESS OR IMPLEA REGRANTED INITY, IMPLIED WARRANTIES, EXPERESS OR IMPLEA REPRESS OR IMPLEAD IN INFINITY ("INFINITY PROPOSE"). AND ANY PARTICULAR VARRANTIES, EXPERESS OR IMPLEAD IN INFINITY PROPOSED, OR NON INFINITY PROPOSED IN INFINITY PROPOSED I