

# Mouse IL-23 Recombinant Protein, eBioscience™

Product Details	
Size	10 µg
Species	Mouse
Expression System	Insect
Amino acid sequence	Heterodimeric, mouse p40, amino acids Met1-Ser355 (Accession # NM_008352), and mouse p19, amino acids Met1-Ala196 (Accession # AF301619)
Molecular weight	55.6 kDa
Class	Recombinant
Type	Protein
Purity	>98% by SDS-PAGE
Endotoxin concentration	<0.01 ng/µg
Activity	ED50 < 625 pg/mL; as measured by the induction of IL-17A in mouse splenocytes.
Conjugate	Unconjugated
Form	Liquid
Concentration	0.1 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 6.0, with 1% BSA
Contains	no preservative
Storage conditions	-80°C

Applications	Tested Dilution	Publications
Western Blot (WB)	Assay-dependent	-
ELISA (ELISA)	Assay-Dependent	-
Functional Assay (FN)	Assay-Dependent	-
Control (Ctrl)	Assay-Dependent	-
Miscellaneous PubMed (Misc)	-	9 Publications

## Product Specific Information

Description: IL-23 is a heterodimeric cytokine composed of the p40 subunit of IL-12 disulfide-linked with a protein p19. p19, like p35 of IL-12, is biologically inactive by itself. IL-23 interacts with IL-12Rbeta1 and an additional, novel beta2-like receptor subunit with STAT4 binding domain, termed IL-23R. IL-23 is secreted by activated mouse and human dendritic cells. Biological activities of mouse IL-23 are distinct from those of mouse IL-12. Mouse IL-23 was found not to induce significant amounts of IFN-gamma. Mouse IL-23 does induce strong proliferation of memory T cells (but not naive T cells), whereas IL-12 has no effect on memory cells. Additionally, mouse IL-23 (but not IL-12) can activate mouse memory T cells to produce the proinflammatory cytokine IL-17. Human IL-23 has biological properties which are less distinct from human IL-12; human IL-23 induces proliferation of memory T cells and induces moderate levels of IFN-gamma production by naive and memory T cells, as compared to IL-12.

Applications Reported: Recombinant mouse IL-23 is biologically active.

Applications Tested: The ED50 of this protein, as measured by IL-17A induction in mouse splenocytes, is less than or equal to 625 pg/mL. This corresponds to a specific activity of greater than  $1.5 \times 10^6$  Units/mg.

Source: Insect cells infected with baculovirus: mouse p40, amino acids met 1-Ser 355, accession # NM\_008352 was co-expressed with mouse p19, amino acids met 1-Ala 196, accession # AF301619.

Bioactivity: The ED50 of this protein, as measured by IL-17A induction in mouse splenocytes, is less than or equal to 625 pg/mL. This corresponds to a specific activity of greater than  $1.5 \times 10^6$  Units/mg.

Endotoxin: Less than 0.01 ng/ug cytokine as determined by the LAL assay. Purity: >98% as determined by SDS-PAGE.

Molecular Weight: The heterodimer of p40, amino acids met23-ser335, cystine-linked to p19, amino acids leu 20-ala 196, has a predicted molecular mass of 55,616. On non-reducing SDS-PAGE the heterodimeric cystine-linked protein migrates as a 60 kDa protein. The DTT reduced protein migrates as 43 kDa and 21 kDa polypeptides.


Storage and handling: For best recovery, quick-spin vial prior to opening. Use in a sterile environment.

Purity: Greater than 90%, as determined by SDS-PAGE.

Aggregation: Less than 10%, as determined by HPLC.

Filtration: 0.2  $\mu$ m post-manufacturing filtered.

### Product Images For Mouse IL-23 Recombinant Protein, eBioscience™



#### Mouse IL-23 Protein (14-8231-63) in WB

Under non-reducing conditions on SDS-PAGE, Mouse IL-23 Recombinant Protein migrates as a 60 kDa protein (Lane NR). The DTT-reduced Mouse IL-23 Recombinant Protein migrates as 43 kDa and 21 kDa polypeptides (Lane R).

## 9 References

### Miscellaneous PubMed (9)

Frontiers in immunology

#### **ATF3 Sustains IL-22-Induced STAT3 Phosphorylation to Maintain Mucosal Immunity Through Inhibiting Phosphatases.**

Authors: Glal D,Sudhakar JN,Lu HH,Liu MC,Chiang HY,Liu YC,Cheng CF,Shui JW

**Species**  
Not Applicable

**Dilution**  
Not Cited

**Year**  
2019

Cellular and molecular gastroenterology and hepatology

#### **Toll-Like Receptor 7 Agonist-Induced Dermatitis Causes Severe Dextran Sulfate Sodium Colitis by Altering the Gut Microbiome and Immune Cells.**

Authors: Kiyohara H,Sujino T,Teratani T,Miyamoto K,Arai MM,Nomura E,Harada Y,Aoki R,Koda Y,Mikami Y,Mizuno S,Naganuma M,Hisamatsu T,Kanai T

**Species**  
Not Applicable

**Dilution**  
Not Cited

**Year**  
2019

[View more Misc references on thermofisher.com](#)

### More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.