

Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 647


Product Details	
Size	1 mg
Species Reactivity	Rabbit
Published Species	Rabbit
Host/Isotope	Goat / IgG
Class	Polyclonal
Type	Secondary Antibody
Conjugate	Alexa Fluor® Plus 647
Immunogen	Gamma Immunoglobins Heavy and Light chains
Form	Liquid
Concentration	2 mg/mL
Purification	Affinity chromatography
Storage buffer	proprietary buffer, pH 6.5
Contains	0.016% Methylisothiazolone, 0.016% Bromonitrodioxane
Storage Conditions	4° C, store in dark
RRID	AB_2633282

Applications	Tested	Dilution	Published
Immunohistochemistry (IHC)	-	1:1000	3 Publications
Miscellaneous PubMed (MISC)	-		3 Publications
Immunocytochemistry (ICC)	✓	1-10 µg/mL	6 Publications
Immunofluorescence (IF)	✓	1-10 µg/mL	6 Publications
Western Blot (WB)	✓	0.05-0.2 µg/mL	3 Publications
Flow Cytometry (Flow)	-	1:1000	1 Publication
Immunohistochemistry - Free Floating (IHC (Free))	-	1:500	1 Publication


Product Specific Information

To minimize cross-reactivity, the goat anti-rabbit IgG whole antibodies have been pre cross-adsorbed against bovine IgG, goat IgG, mouse IgG, rat IgG, and human IgG. Cross-adsorption or pre-adsorption is a purification step to increase specificity of the antibody resulting in less background staining and cross-reactivity. The secondary antibody solution is passed through a column matrix containing immobilized serum proteins from potentially cross-reactive species. Only the nonspecific-binding secondary antibodies are captured in the column, and the highly specific secondaries flow through. Further passages through additional columns result in highly cross-adsorbed preparations of secondary antibody. The benefits of these extra steps are apparent in multiplexing/multicolor-staining experiments where there is potential cross-reactivity with other primary antibodies or in tissue/cell fluorescent staining experiments where there may be the presence of endogenous immunoglobulins.


Using conjugate solutions: Centrifuge the protein conjugate solution briefly in a microcentrifuge before use; add only the supernatant to the experiment. This step will help eliminate any protein aggregates that may have formed during storage, thereby reducing nonspecific background staining. Because staining protocols vary with application, the appropriate dilution of antibody should be determined empirically.



Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A32733) in IF
Immunofluorescent analysis of postsynaptic density protein (PSD-95) in E18 Sparague Dawley primary cortical neuronal cells. The cells were fixed with 4% formaldehyde for 15 mins, permeabilized with 0.25% Triton X-100 in PBS for 10 mins, and blocked with 3% BSA in PBS for 30 mins at RT. Cells were stained with a PSD-95 rabbit polyclonal antibody (Product # 51-6900) at a dilution of 1:200 in 3% BSA in PBS for 1 hr at RT, and then incubated with Invitrogen Alexa Fluor Plus 647 goat anti-rabbit IgG secondary antibody (Product # A32733) at a dilution of 1:1000 for 1 hr at RT. Nuclei were stained with Hoechst 33342 (product # H3570). The image contains overlay of neuronal PSD-95 (far red) and nuclei (blue). Images were taken on a Zeiss LSM 710 confocal microscope at 40X magnification.



Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A32733) in IF
Immunofluorescent analysis of ZO-1 in A549 cells. The cells were fixed with 4% formaldehyde for 15 mins, permeabilized with 0.25% Triton X-100 in PBS for 10 mins, and blocked with 3% BSA in PBS for 30 mins at RT. Cells were stained with a ZO-1 rabbit antibody (Product # 40-2300) at a dilution of 1:200 in 3% BSA in PBS for 1 hr at RT, and then incubated with Invitrogen Alexa Fluor Plus 647 goat anti-rabbit IgG secondary antibody (Product # A32733) at a dilution of 1:1000 for 1 hr at RT. Nuclei were stained with Hoechst 33342 (Product # H3570). The image contains overlay of ZO-1 (far red) and nuclei (blue). Images were taken on a Zeiss LSM 710 confocal microscope at 40X magnification.



Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A32733) in IF
Immunofluorescent analysis of PMP70 in A549 cells. The cells were fixed with 4% formaldehyde for 15 mins, permeabilized with 0.25% Triton X-100 in PBS for 10 mins, and blocked with 3% BSA in PBS for 30 mins at RT. Cells were stained with a PMP70 rabbit antibody at a dilution of 1:200 in 3% BSA in PBS for 1 hr at RT, and then incubated with Invitrogen Alexa Fluor Plus 647 goat anti-rabbit IgG secondary antibody (Product # A32733) at a dilution of 1:1000 for 1 hr at RT. Nuclei were stained with Hoechst 33342 (product # H3570). The image contains overlay of PMP70 (far red) and nuclei (blue). Images were taken on a Zeiss LSM 710 confocal microscope at 40X magnification.

[View more figures on thermofisher.com](https://www.thermofisher.com)

23 References

Immunohistochemistry (3)

Frontiers in cellular neuroscience

Region-Specific Reductions in Morphometric Properties and Synaptic Colocalization of Astrocytes Following Cocaine Self-Administration and Extinction.

Authors: Testen A, Sepulveda-Orengo MT, Gaines CH, Reissner KJ

Species

Rabbit
Not Applicable

Dilution

1:1000
Not Cited

Year

2018

Nature communications

Precise inhibitory microcircuit assembly of developmentally related neocortical interneurons in clusters.

Authors: Zhang XJ, Li Z, Han Z, Sultan KT, Huang K, Shi SH

Species

Not Applicable

Dilution

1:1000

Year

2017

[View more IHC references on thermofisher.com](#)

Miscellaneous PubMed (3)

Frontiers in cellular neuroscience

Region-Specific Reductions in Morphometric Properties and Synaptic Colocalization of Astrocytes Following Cocaine Self-Administration and Extinction.

Authors: Testen A, Sepulveda-Orengo MT, Gaines CH, Reissner KJ

Species

Rabbit
Not Applicable

Dilution

1:1000
Not Cited

Year

2018

Nature

A semi-synthetic organism that stores and retrieves increased genetic information.

Authors: Zhang Y, Ptacin JL, Fischer EC, Aerni HR, Caffaro CE, San Jose K, Feldman AW, Turner CR, Romesberg FE

Species

Rabbit
Not Applicable

Dilution

1:20,000
Not Cited

Year

2017

[View more MISC references on thermofisher.com](#)

More applications with references on thermofisher.com

ICC (6)

IF (6)

WB (3)

Flow (1)

IHC (Free) (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.