



Chicken anti-Rat IgG (H+L) Secondary Antibody, TRITC


| Product Details | | |
|----------------------------|---|--|
| Size | 1 mg | |
| Species Reactivity | Rat | |
| Host/Isotype | Chicken / IgY | |
| Class | Polyclonal | |
| Туре | Secondary Antibody | |
| Conjugate | TRITC | |
| Excitation/Emission Max | 552/578 nm | |
| Immunogen | Rat IgG whole molecule | |
| Form | Liquid | |
| Concentration | 1 mg/mL | |
| Purification | IgG fraction | |
| Storage buffer | 0.02M potassium phosphate, pH 7.2, with 0.15M NaCl, 10mg/mL BSA | |
| Contains | 0.01% sodium azide | |
| Storage conditions | Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles. Store in the dark. | |
| RRID | AB_10984821 | |

| Applications | Tested Dilution | Publications |
|------------------------------|-----------------|--------------|
| Immunocytochemistry (ICC/IF) | 1:1,000-1:5,000 | - |
| Flow Cytometry (Flow) | 1:500-1:2500 | - |

Product Specific Information

Store product as a concentrated solution. Centrifuge briefly prior to opening the vial.

Product Images For Chicken anti-Rat IgG (H+L) Secondary Antibody, TRITC



Support (US): 1 800 955 6288

E-mail: CustomerCare@thermofisher.com

Rat IgG (H+L) Secondary Antibody (PA1-29949) in ICC/IF

Immunofluorescence analysis of Chicken anti-Rat IgG Secondary Antibody, TRITC conjugate was performed using A549 cells stained with alpha Tubulin (YL1 /2) Rat Monoclonal Antibody (Product # MA1-80017). The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 10 minutes, blocked with 1% BSA for 1 hour and labeled with 2µg/mL Rat primary antibody for 3 hours at room temperature. Chicken anti-Rat IgG Secondary Antibody, TRITC conjugate (Product # PA1-29949) was used at a concentration of 2µg/mL in phosphate buffered saline containing 0.2 % BSA for 45 minutes at room temperature, for detection of alpha Tubulin in the cytoplasm (Panel a: red). Nuclei (Panel b: blue) were stained with DAPI in SlowFade® Gold Antifade Mountant (Product # S36938). F-actin was stained with Alexa Fluor® 488 Phalloidin (Product # A12379), 1:300) (Panel c: green). Panel d represents the composite image. No nonspecific staining was observed with the secondary antibody alone (panel f), or with an isotype control (panel e). The images were captured at 60X magnification.

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation For Research 109 of Only, motive use in inaginastic procedures, Not. on the agent and intermediate and inter REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals

2