





CD20 Monoclonal Antibody (2H7), APC

MA1-82679 Product data sheet **Catalog Number**

Details		Species Reactivity	
Size	100 Tests	Species reactivity	Н
Host/Isotope	Mouse / IgG2b	Published species	Н
Class	Monoclonal	Tested Applications	D
Туре	Antibody	Flow Cytometry (Flow)	N
Clone	2H7	Published Applications	
Immunogen	Proprietary Immunogen.	Flow Cytometry (Flow)	S
Conjugate	APC	* Suggested working dilutions are given as a guide only. It is recommend experiment using appropriate negative and positive controls.	
Form	Lyophilized		
Concentration	Conc. Not Determined		
Purification	Protein A		
Storage buffer	PBS, pH 7.4, with 1% BSA, 5% sucrose		
Contains	0.09% sodium azide		
Storage Conditions	4° C, do not freeze		

Species Reactivity			
Species reactivity	Human, Rhesus monkey		
Published species	Human		
Tested Applications	Dilution *		
Flow Cytometry (Flow)	Neat		
Published Applications			
Flow Cytometry (Flow)	See 1 publications below		
Published Applications			

ed that the user titrate the product for use in their own

Product specific information

Reconstitute with 1 mL of distilled water. After reconstitution, store product undiluted at 4°C. For FACS analysis, use 10 µL of the suggested working dilution to label 1x10⁶ cells in 100 μL. Mouse anti Human CD20 antibody, clone 2H7 recognizes the human CD20 cell surface antigen, a 33-37 kDa non-glycosylated phosphoprotein.

Background/Target Information

CD20 is a non-glycosylated surface phosphoprotein that has a molecular weight range of 33-37 kDa depending on the degree of phosphorylation. CD20 is expressed on mature and most malignant B cells, in a subpopulation of T lymphocytes and follicular dendritic cells. CD20 expression on B cells is synchronous with the expression of surface IgM and it regulates transmembrane calcium conductance, cell cycle progression and B-cell proliferation. CD20 is also associated with lipid rafts, but the intensity of this association depends on extracellular triggering, employing CD20 conformational change, and/or BCR (B cell antigen receptor) aggregation. After the receptor ligation, BCR and CD20 colocalize and then rapidly dissociate before BCR endocytosis, whereas CD20 remains at the cell surface. CD20 serves as a useful target for antibody-mediated therapeutic depletion of B cells, as it is expressed at high levels on most B-cell malignancies, but does not become internalized or shed from the plasma membrane following monoclonal antibody treatment. Diseases associated with CD20 dysfunction include Ms4a1-related common variable immune deficiency.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.

PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PR FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICI for research only and is not to be used for any other purpose, including without limitation, unauthorized cor



PubMed References For CD20 Monoclonal Antibody (2H7), APC 1 Flow Cytometry References Species / Dilution Summary MA182679 was used in flow cytometry to correlate biomarkers associated with coagulation, monocyte activation, and inflammation in elite controllers of HIV infection Human / Not Cited The Journal of infectious diseases (2014; 209: 931) "Evidence for innate immune system activation in HIV type 1-infected elite controllers." Author(s):Krishnan S,Wilson EM,Sheikh V,Rupert A,Mendoza D,Yang J,Lempicki R,Migueles SA,Sereti I PubMed Article URL:http://dx.doi.org/10.1093/infdis/jit581

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACE OR REFUND FOR THE RESULT OF (10) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (10) MISUSE, FAULT OR REGILGENCE OF OR BY BUYER, (11) USE or THE PRODUCTS IN A MANNIER FOR WHICH THEY WERE NOT DESIGNED, (10) (1) MIPOPOPE STORAGE AND HANDLING OF THE PRODUCTS. In Library Expressed stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.

