





RGS14 Polyclonal Antibody

Catalog Number PA5-30396 Product data sheet

Details	
Size	100 μL
Host/Isotope	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Immunogen	Recombinant fragment corresponding to a region within amino acids 286 and 558 of Human RGS14
Conjugate	Unconjugated
Form	Liquid
Concentration	0.79 mg/mL
Purification	Antigen affinity chromatography
Storage buffer	0.1M tris glycine, pH 7, with 20% glycerol
Contains	0.01% thimerosal
Storage Conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

Species Reactivity		
Species reactivity	Human	
Tastad Applications	Dilution t	
Tested Applications	Dilution *	
Tested Applications Western Blot (WB)	Dilution * 1:500-1:3,000	

^{*} Suggested working dilutions are given as a guide only. It is recommended that the user titrate the product for use in their own experiment using appropriate negative and positive controls.

Product specific information

Recommended positive controls: Raji. Predicted reactivity: Mouse (82%), Rat (80%), Bovine (84%). Store product as a concentrated solution. Centrifuge briefly prior to opening the vial.

Background/Target Information

RGS14 is a member of the regulator of G-protein signaling family which contains one RGS domain, two Raf-like Ras-binding domains (RBDs), and one GoLoco domain. RGS14 attenuates the signaling activity of G-proteins by binding, through its GoLoco domain, to specific types of activated, GTP-bound G alpha subunits. Acting as a GTPase activating protein (GAP), the protein increases the rate of conversion of the GTP to GDP. The resulting hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. Alternate transcriptional splice variants of RGS14 have been observed but have not been thoroughly characterized. Further, RGS14 acts as a positive modulator of microtubule polymerisation and spindle organization through a G(i)-alpha-dependent mechanism; plays a role in cell division; is required for the nerve growth factor (NGF)-mediated neurite outgrowth, and is involved in stress resistance. RGS14 protein may also be involved in visual memory processing capacity and hippocampal-based learning and memory.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT, BUYER'S EXCLUSINE REALED FOR ROCKETOR POR REFUND FOR PORTOR PORT OF REFUND FOR PORTOR PORTO



Product Images For RGS14 Polyclonal Antibody

KDa 170 — A 130 — 95 — 72 — ← 55 —

RGS14 Antibody (PA5-30396) in WB

Western Blot using RGS14 Polyclonal Antibody (Product # PA5-30396). Sample (30 µg of whole cell lysate). Lane A: Raji . 7.5% SDS PAGE. RGS14 Polyclonal Antibody (Product # PA5-30396) diluted at 1:1,000.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization

Products are warranted to operate or perform substantially in continemance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package insents ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACE OR REFUND FOR THE RESULT OF (10) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (10) MISUSE, FAULT OR REGILGENCE OF OR BY BUYER, (11) USE or THE PRODUCTS IN A MANNIER FOR WHICH THEY WERE NOT DESIGNED, (10) (1) MIPOPOPE STORAGE AND HANDLING OF THE PRODUCTS. In Library Expressed stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.

