





RNF216 Polyclonal Antibody

Catalog Number PA1-41317 Product data sheet

Details	
Size	100 μg
Host/Isotope	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Immunogen	Synthetic peptide corresponding to residues 350-400 of human TRIAD3A.
Conjugate	Unconjugated
Form	Liquid
Concentration	0.5 mg/mL
Purification	Protein G
Storage buffer	PBS with 0.2% gelatin
Contains	0.05% sodium azide
Storage Conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

Human
Dilution *
1-3 μg/mL

^{*} Suggested working dilutions are given as a guide only. It is recommended that the user titrate the product for use in their own experiment using appropriate negative and positive controls.

Product specific information

Suggested positive control: human spleen protein, antigen standard for RNF216 (transient overexpression lysate), human lung protein, human ovary protein, human pancreas protein, human skeletal muscle protein, human heart protein, human kidney protein, human liver protein, human brain protein, human brain tissue, heart tissue, kidney tissue, liver tissue, lung tissue, ovary tissue, pancreas tissue, small intestine tissue, skeletal muscle tissue, spleen tissue, human small intestine protein.

Background/Target Information

Activation of NF-B as a result of Toll-like receptor (TLR) and IL-1 receptor signaling is a major component of innate immune responses. Signals from these receptors are relayed by a number of adapter molecules such as TRIF, TIRAP, and MyD88. Several regulatory mechanisms exist to control TLR signal transduction, including the inhibition of TLR expression and signaling by molecules such as ST2 and SIGIRR. Another mechanism is by the ubiquitinization of selected TLRs by TRIAD3A, an E3 ubiquitin-protein ligase. TRIAD3A is a RING finger protein that can bind to TLR4 and TLR9, and to a lesser extent TLR3 and TLR5, catalyzing the ubiquitization of these molecules. Overexpression of TRIAD3A promoted the nearly complete degradation of TLR4 and TLR9; this reduction was reflected in the decreased signal-specific activation by ligands specific for these TLRs. Conversely, depletion of TRIAD3A resulted in enhanced TLR activation.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.


For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT, BUYER'S EXCLUSINE REALED FOR ROCKETOR POR REFUND FOR PORTOR PORT OF REFUND FOR PORTOR PORTO



Product Images For RNF216 Polyclonal Antibody



RNF216 Antibody (PA1-41317) in WB

Western blot analysis of RNF216 in human (A) Brain; (B) Heart; (C) Small Intestine; (D) Kidney; (E) Liver; (F) Lung; (G) Skeletal Muscle; (H) Pancreas; (I) Spleen; (J) Ovary; (K) Testis. Samples were incubated in RNF216 polyclonal antibody (Product # PA1-41317) using a dilution of 1 μg/mL.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACE OR REFUND FOR THE RESULT OF (10) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (10) MISUSE, FAULT OR REGILGENCE OF OR BY BUYER, (11) USE or THE PRODUCTS IN A MANNIER FOR WHICH THEY WERE NOT DESIGNED, (10) (1) MIPOPOPE STORAGE AND HANDLING OF THE PRODUCTS. In Library Expressed stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.

